

The Istanbul consensus update: a revised ESHRE/ALPHA consensus on oocyte and embryo static and dynamic morphological assessment^{†,‡}

The Working Group on the update of the ESHRE/ALPHA Istanbul Consensus, Giovanni Coticchio ^{1,*}, Aisling Ahlström ², Gemma Arroyo³, Basak Balaban⁴, Alison Campbell ^{5,6}, Maria José De Los Santos ^{7,8}, Thomas Ebner ⁹, David K. Gardner ^{10,11}, Borut Kovačič ¹², Kersti Lundin¹³, M. Cristina Magli ¹⁴, Saria Mcheik ¹⁵, Dean E. Morbeck ^{16,17}, Laura Rienzi¹⁸, Ioannis Sfontouris ¹⁹, Nathalie Vermeulen ¹⁵, and Mina Alikani ^{20,*}

¹IVIRMA Global Research Alliance, IVIRMA Italia, Rome, Italy

²IVIRMA Global Research Alliance, LIVIO, Göteborg, Sweden

³Dpt d'Obstetricia i Ginecologia, Institut Universitari Dexeus, Barcelona, Spain

⁴Assisted Reproduction Unit, VKF American Hospital of Istanbul, Istanbul, Turkiye

⁵CARE Fertility Group, Nottingham, UK

⁶University of Kent, Kent, UK

⁷IVIRMA Valencia Global Research Alliance, IVF Laboratory, Valencia, Spain

⁸Fundación IVI Instituto de Investigaciones Sanitarias, Valencia, Spain

⁹Gynecology Obstetrics and Gynecological Endocrinology, Kepler Universitätsklinikum GmbH, Linz, Austria

¹⁰Melbourne IVF, East Melbourne, VIC, Australia

¹¹School of BioSciences, University of Melbourne, Parkville, VIC, Australia

¹²Department for Reproductive Medicine and Gynecological Endocrinology, University Medical Centre Maribor, Maribor, Slovenia

¹³Dept of Obstetrics and Gynecology, The Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

¹⁴SISMER Reproductive Medicine Unit, Bologna, Italy

¹⁵ESHRE, Strombeek-Bever, Belgium

¹⁶Genea Fertility, Sydney, NSW, Australia

¹⁷Department of Obstetrics and Gynecology, Monash University, Melbourne, VIC, Australia

¹⁸IVIRMA Global Research Alliance, Rome, Italy

¹⁹Hygeia IVF Embryogenesis, Athens, Greece

²⁰Alpha Scientists in Reproductive Medicine, London, UK

*Correspondence address. IVIRMA Global Research Alliance, IVIRMA ITALIA, Via dei Monti Parioli 6, 00197 Roma, Italy. E-mail: giovanni.coticchio@gmail.com

 <http://orcid.org/0000-0003-1635-9205> (G.C.); Alpha Scientists in Reproductive Medicine, Suite 213 Boundary House, Boston Road, London, W7 2QE, UK.

E-mail: mina.alikani@embryos.net guidelines@eshre.eu  <http://orcid.org/0000-0001-6140-0321> (M.A.)

[†]ESHRE Pages content is not externally peer reviewed. The manuscript has been approved by the Executive Committee of ESHRE.

[‡]This article has been simultaneously co-published, with permission, in Human Reproduction, and Reproductive Biomedicine Online.

ABSTRACT

STUDY QUESTION: What are the current recommended criteria for morphological assessment of oocytes, zygotes, and embryos?

SUMMARY ANSWER: The present ESHRE/Alpha Scientists in Reproductive Medicine consensus document provides several novel recommendations to assess oocyte and embryo morphology and rank embryos for transfer.

WHAT IS KNOWN ALREADY: A previous Alpha Scientists in Reproductive Medicine/ESHRE consensus on oocyte and embryo morphological assessment was published in 2011. After more than a decade, and the integration of time-lapse technology into embryo culture and assessment, a thorough review and update was needed.

STUDY DESIGN, SIZE, DURATION: A working group consisting of Alpha Scientists in Reproductive Medicine executive committee members and ESHRE Special interest group of Embryology members formulated recommendations on oocyte and embryo assessment.

PARTICIPANTS/MATERIALS, SETTING, METHODS: The working group included 17 internationally recognized experts with extensive experience in clinical embryology. Seven members represented Alpha Scientists in Reproductive Medicine and eight members represented ESHRE, along with two methodological experts from the ESHRE central office. Based on a systematic literature search and discussion of existing evidence, the recommendations of the Istanbul Consensus (2011) were reassessed and, where appropriate, updated based on consensus within the working group. A stakeholder review was organized after the updated draft was finalized. The final version was approved by the working group, the Alpha executive committee and the ESHRE Executive Committee.

Received: December 13, 2024. Editorial decision: January 17, 2025.

© The Author(s) 2025. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

MAIN RESULTS AND THE ROLE OF CHANCE: This updated consensus paper provides 20 recommendations focused on the timeline of preimplantation developmental events and morphological criteria for oocyte, zygote, and embryo assessment. Based on duration of embryo culture, recommendations are given on the frequency and timing of assessments to ensure consistency and effectiveness.

LIMITATIONS, REASONS FOR CAUTION: Several criteria relevant to oocyte and embryo morphology have not been well studied, leading to either a recommendation against their use for grading or for their use in ranking rather than grading. Future updates may require further revision of these recommendations.

WIDER IMPLICATIONS OF THE FINDINGS: This document provides embryologists with advice on best practices when assessing oocyte and embryo quality based on the most recent evidence.

STUDY FUNDING/COMPETING INTEREST(S): The consensus meeting and writing of the paper were supported by funds from ESHRE and Alpha Scientists in Reproductive Medicine. The working group members did not receive any payment. G.C. declared payments or honoraria for lectures from Gedeon Richter and Cooper Surgical. A.C. declared text book royalties (Mastering Clinical Embryology, published 2024), consulting fees from Cooper Surgical, Gedeon Richter and TMRW Life Sciences, honoraria for lectures from Merck, Ferring, and Gedeon Richter, and participation in the HFEA Scientific Advances Committee; she also disclosed being treasurer and vice-president of Alpha Scientists in Reproductive Medicine, a shareholder in Care Fertility Limited and Fertile Mind Limited, and having stock options in TMRW Life Sciences and U-Ploid Biotechnology Ltd. L.R. declared consulting fees from Organon, payments or honoraria for lectures from Merck, Organon, IBSA, Finox, Geden Richter, Origio, Organon, Ferring, Foundation IVI; she also disclosed being a member of the Advisory Scientific Board of IVIRMA (Paid) and a member of the Advisory Scientific Board of Nterilizer (unpaid). I.S. declared payments or honoraria for lectures from Vitrolife and Cooper Surgical, and stock options from Alife Health. M.A. declared payments or honoraria for lectures from Vitrolife and support for attending meetings from Vitrolife and Cooper Surgical (both unrelated to this manuscript). The other authors have no conflicts of interest to declare.

DISCLAIMER: This Good Practice Recommendations (GPRs) document represents the consensus views of the members of this working group based on the scientific evidence available at the time of the meeting. GPRs should be used for information and educational purposes. They should not be interpreted as setting a standard of care or be deemed inclusive of all proper methods of care or be exclusive of other methods of care reasonably directed to obtaining the same results. They do not replace the need for application of clinical judgement to each individual presentation, or variations based on locality and facility type.

Keywords: embryo / oocyte / cleavage stage embryo / morula / blastocyst / morphology / morphokinetics / time-lapse

Introduction

Assessment of human embryo development is an essential, but challenging, task in the IVF laboratory. Embryos are assessed by embryologists to select the most likely to be viable for intrauterine transfer, cryopreservation or biopsy for preimplantation genetic testing (PGT). Since the early days of IVF in the 1980s when embryos were optimistically viewed as ‘nice, very nice, or very very nice’ (Jacques Cohen, personal communication), a relatively large number of early embryo morphological features have been identified and investigated for their association with viability, implantation, live birth and chromosomal status. Yet, morphology assessment remains largely subjective and prone to inter- and intra-observer and inter-laboratory variability (Arce et al., 2006; Baxter Bendus et al., 2006; Martínez-Granados et al., 2017; Storr et al., 2017).

In the past decade, the most significant advancement in embryo assessment has been the introduction of time-lapse microscopy technologies (TLT). This has led to the emergence of ‘morphokinetics’. As the term implies, morphokinetics represents the integration of morphology (the form and structure of embryos) with kinetics (the dynamics of their development), providing a comprehensive framework for understanding and evaluating embryo development in vitro. These technologies allow continuous observation of embryo development, with minimal manipulation or perturbation of culture (ESHRE Working group on Time-lapse technology et al., 2020).

Hundreds of papers have been published on embryo assessment. The studies are mostly retrospective and heterogeneous with respect to some key parameters including patient population, outcome measures, control for confounders, laboratory procedures, and embryo culture conditions. Furthermore, morphokinetic studies, as well as classical morphological studies, may be influenced by maternal age, smoking status, ovarian stimulation protocols, and insemination methods, among other factors (Braga et al., 2015; Ubaldi et al., 2016; Grøndahl et al., 2017;

Barrie et al., 2021a; Bamford et al., 2022). Nonetheless, TLT observations have significantly contributed to our understanding of developmental events, and morphology assessments are now enhanced by morphokinetics.

Over a decade ago, Alpha Scientists in Reproductive Medicine (ALPHA) and ESHRE special interest group of Embryology collaborated to produce the Istanbul Consensus on assessing oocytes, zygotes, and embryos (Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology, 2011).

The Istanbul Consensus (2011) established common criteria and terminology for grading oocytes, zygotes and embryos, which are now updated in this paper through close examination, compilation, analysis and interpretation of data published in the intervening years. Most importantly, the new recommendations incorporate some embryo morphokinetic features that have been elucidated since the introduction of TLT and that can inform and complement the static observation approach. The aim of this document is to help re-establish standard terminology and assessment criteria across laboratories.

Terminology

Embryologists routinely make decisions on disposition of oocytes and embryos, that is, whether they are clinically usable or should be discarded. Clinical use of oocytes and embryos is defined as use for an IVF/ICSI treatment, biopsy/PGT, cryopreservation, transfer, and donation.

In the updated set of recommendations provided in this manuscript, the working group used the terms embryo grading, ranking, and selection. *Embryo grading* is the evaluation of embryos using a specific set of criteria to assign a quality score: the number, size, and shape of blastomeres, the degree of fragmentation, the inner cell mass (ICM) and trophectoderm (TE) morphology and expansion, etc. *Embryo ranking* refers to the procedure of prioritizing clinically usable embryos based on grading and other assessment criteria, from most to least favourable for transfer. Embryos are ranked according to their estimated potential for

implantation and development, which is determined by morphological and, when available, genetic factors. This is a prioritization of which embryo(s) to transfer first. Embryo selection for transfer involves consideration of ranking and other factors to select embryos for transfer into the uterus. The goal is to select the embryo(s) with the highest likelihood of resulting in a successful pregnancy and live birth.

Materials and methods

The present good practice recommendations document is the result of a multiple virtual meetings over a 1-year period and a 2-day consensus meeting of a working group (WG) of expert professionals representing ALPHA and ESHRE. As a starting point for the update process, a survey was created to collect information on current practice in ART centres regarding the application of the Istanbul Consensus (2011) recommendations. The questionnaire had three sections, with mostly multiple-choice answers; it inquired about the country of practice, the classification system in use, the adoption of the Istanbul Consensus (2011) recommendations, and considerations regarding the use of other technologies including TLT, artificial intelligence (AI), and PGT (Supplementary Data SI). Respondents were ensured anonymity as no identifying information was requested. Nonetheless, they were not allowed to take the survey more than once from the same device. The survey was distributed among ALPHA and ESHRE members and posted on the two societies' websites and social media pages. It was requested that one senior representative of the centre complete the survey. In total, 833 responses were collected between 21 November 2022 and the second of January 2023. Survey results can be found in Supplementary Data SII.

In addition, data on oocyte and embryo static and dynamic assessment published up to May 2024 were collected from the literature in PubMed/MEDLINE. All titles and abstracts were screened. Only papers considered to be relevant were selected and included in the text. Papers published after this date were manually included if deemed relevant for this manuscript. References retrieved from the literature were complemented with further key references identified by the WG members. The paper quality was assessed using the GRADE Pro software (McMaster University, USA). The recommendations for clinical practice were formulated based on the expert opinion of the WG, taking into consideration the available evidence and results of the survey.

During the consensus meeting, the results of the survey, scientific evidence and personal clinical experience were integrated into presentations by experts on specific topics. After the presentation of the topic, each proposed recommendation for assessment was discussed until consensus was reached within the group. An updated text including the most relevant papers was prepared and consensus points were included. After approval of the manuscript by the meeting participants, the final draft was published on the ALPHA and ESHRE websites between 17 May 2024 and 17 June 2024 for stakeholder review. In total, 157 comments were received and considered when relevant. The review report is available on www.eshre.eu/guidelines and <https://alphascientists.org/>.

The final draft of this manuscript was approved by the executive committee members of both societies. Abbreviations used throughout this article are listed in Supplementary Data SIII.

Current data on oocyte and embryo assessment criteria

1. Expected timeline of embryo development

Development of the human embryo begins with fertilization and continues with a series of restrictive mitotic events (cleavage) each of which doubles the cell number as the embryo develops from a single cell into a multicellular blastocyst (Ciray *et al.*, 2014). At fertilization, once the two pronuclei break down, paternal and maternal chromosomes are assembled into a bipolar mitotic spindle, before sister chromatids are orderly segregated in the first two blastomeres at first cleavage. The resulting undifferentiated daughter cells are expected to be genetically identical. In the initial developmental phases, blastomere function is under the primary control of a sophisticated regulatory mechanism guided by maternal factors (Sha *et al.*, 2020). However, recent studies have investigated the fine details of the first event of chromosome segregation in the human embryo, revealing a highly error-prone mechanism (Currie *et al.*, 2022). Although the exact timing is yet to be elucidated, embryonic genome activation is well underway by the 8-cell stage, with the concomitant degradation of maternal transcripts (Braude *et al.*, 1988; Vassena *et al.*, 2011; Asami *et al.*, 2022; Yuan *et al.*, 2023).

Since the competence of the human embryo is also reflected in its developmental timeline, assessment of morphology should be in accordance with predefined times.

The original Istanbul Consensus (2011) on embryo assessment proposed specific timings for observations of fertilized oocytes and embryos, and their expected stage of development at these time points. These timings were relative to the insemination time and aimed to reflect when the events of interest occur generally (Table 1). Times for observations were provided for the following stages: fertilization, syngamy, early cleavage, Day-2, -3, -4 and -5 embryo assessment. The Istanbul Consensus (2011) differentiated between IVF- and ICSI-derived embryos only for one stage of development: early cleavage. Specifically, the 2-cell stage was proposed to be checked 2 h earlier post ICSI (26 ± 1 h post-insemination (hpi)), than IVF (28 ± 1 hpi) (Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology, 2011). The rationale behind this suggestion is that pronuclear formation post IVF is observed about 1 h later than post ICSI (Nagy *et al.*, 1998), where the cumulus-corona complex, zona pellucida (ZP) and oolemma are bypassed, conserving the time required for the spermatozoon to traverse this path (Payne *et al.*, 1997).

Studies have shown that early cleavage is an independent predictor of embryo quality (in terms of cell number and morphology at later cleavage stages), blastocyst formation, pregnancy and birth, although there were apparent differences between IVF- and ICSI-derived embryos (Shoukir *et al.*, 1997; Lundin *et al.*, 2001; Van Montfoort *et al.*, 2004).

Several subsequent reports of the relative morphokinetic timings of IVF- and ICSI-derived embryos have been described in the literature and were considered in this revised version of the Istanbul Consensus. For example, several studies reported that only the timing of the first cleavage was affected by fertilization method, with IVF embryos reaching the 2-cell stage significantly later than their ICSI counterparts (Dal Canto *et al.*, 2012b; Kirkegaard *et al.*, 2016). Another study detected comparative delays in IVF embryo development beyond the 2-cell stage of 1.5 ± 1.1 h (Bodri *et al.*, 2015). A recent randomized controlled study compared morphokinetics of 373 sibling IVF and ICSI embryos and reported that only time to 2-cell (t2) was significantly delayed in IVF embryos (De Munck *et al.*, 2022). A large TLT study

Table 1. Time lapse data generated reference timings related to specific embryo developmental stage assessments.

Istanbul Consensus 2011			2024		
Type of observation	Timing (hpi)	Expected stage of development	Median time to reach developmental stage (rounded to nearest hour)	Assessment time for each developmental stage to give highest chance of observation (hpi). Rounded. After fertilisation check, all ± 1 hour	Proportion expected to be at stage required for specific assessment. Rounded.
Fertilisation check	17 \pm 1	Pronuclear stage	N/A	16-17 (ICSI or IVF)	98% with visible pronuclei (Barrie <i>et al.</i> , 2021b)
Syngamy check	23 \pm 1	Expect 50% to be in syngamy (up to 20% may be at 2 cell stage)	tPNf (time to pronuclear fading) 23 (ICSI) 24 (IVF)	25 (ICSI) 26 (IVF)	53% 53%
Early cleavage check	26 \pm 1 (ICSI) 28 \pm 1 (IVF)	2 cell stage	t2 (time to 2 cell) 26 (ICSI) 27 (IVF)	31 (ICSI) 32 (IVF)	77% 79%
Day-2 embryo assessment	44 \pm 1	4 cell stage	t4 (time to 4 cell) 38 (ICSI) 39 (IVF)	43 (ICSI) 45 (IVF)	64% 67%
Day-3 embryo assessment	68 \pm 1	8 cell stage	t8 (time to 8 cell) 57 (ICSI) 58 (IVF)	63 (ICSI) 65 (IVF)	49% 51%
Day-4 embryo assessment	92 \pm 2	Morula	tM (time to morulae) 89 (ICSI) 91 (IVF)	93 (ICSI) 95 (IVF)	47% 44%
Day-5 embryo assessment	116 \pm 2	Blastocyst	tB (time to full blastocyst) 108 (ICSI) 107 (IVF)	108 (ICSI) 108 (IVF)	47% 52%
			tEB (time to expanded blastocyst) 113 (ICSI) 113 (IVF)	111 (ICSI) 112 (IVF)	34% 34%

Morphokinetic timings are obtained from manually annotated embryos in vitro (n = 140 872 2PNs—56 066 IVF and 84 806 ICSI) (Unpublished Care Fertility multicentre data 2013–2022), fresh oocytes only. Nomenclature and definitions are based on Ciray *et al.* (2014). Regarding Days 6 and 7 observations, this dataset does not have sufficient data available to offer guidance for observation. However, see Section 6 (blastocyst stage) regarding assessment of blastocysts beyond Day 5.

Hpi, hours post-insemination.

of 2376 embryos reported that t2 was 0.98h earlier in ICSI-derived embryos (excluding those from donor sperm), while time to initiation of blastulation (tSB) and time to full blastocyst (tB) were 1.157 and 1.510h later, respectively, compared with IVF-derived embryos (Barrie et al., 2021a).

Furthermore, many morphokinetic-based studies have investigated the possible influence of other intrinsic and extrinsic factors on the timing of embryo development (e.g. BMI, age, culture media and oxygen concentration) (ESHRE Working group on Time-lapse technology et al., 2020). Two of the most studied patient variables are age of gamete providers and BMI albeit with varying findings and no meta-analyses or definitive studies yet available (Lebovitz et al., 2021; Setti et al., 2021; Bellver, 2022; Boucret et al., 2022; Hoek et al., 2022).

Whether ovarian stimulation protocol impacts embryo developmental timing has also been investigated using morphokinetic analyses, with some apparent differences during early cleavage stages, but no effect on overall embryo quality (Barrie et al., 2017a; Mumusoglu et al., 2017; Dietrich et al., 2020).

Other factors known to affect embryo development, such as temperature and pH, can influence embryo morphokinetics; lower temperature and culture medium pH drift (typically in an alkaline direction) are associated with slower embryo development (Swain, 2015; Wale and Gardner, 2016). The impact of oxygen level during culture, a major influencer of embryo development, has not been extensively studied. However, development and implantation rates decrease when atmospheric oxygen level is employed, compared with lower, more physiological levels (Quinn and Harlow, 1978; Gardner and Kelley, 2017). Using TLT imaging, and similar to data in the mouse (Wale and Gardner, 2010), a prospective study compared the developmental timings of embryos according to oxygen tensions, reporting significantly slower development in embryos cultured in 20% oxygen compared with 5% (Kirkegaard et al., 2013).

There has been discussion regarding possible unconscious bias in selection of faster developing embryos, which may impact the sex ratio. However, a recent large study showed sex ratios, from an IVF program using algorithmic morphokinetic selection, to be in line with the World Health Organisation's (WHO) reported secondary sex ratios for natural conception (Smith et al., 2024).

Utilizing TLT, a number of heterogeneous studies have compared developmental timings according to the chromosomal sex of the embryo, with conflicting results (e.g. Bodri et al., 2016; Serdarogullari et al., 2014). More comparative large studies are needed, however, Fraire-Zamora et al. (2023) aimed to avoid confounding factors by using strict inclusion criteria and reported no significant differences in morphokinetics between male and female embryos (Fraire-Zamora et al., 2023).

Another area of scrutiny has been embryo chromosome status. A recent systematic review and meta-analysis incorporating over 40 000 embryos concluded that ten morphokinetic variables were significantly delayed in aneuploid embryos, most notably from t8 (development at the 8-cell stage) to the expanded blastocyst stage (Bamford et al., 2022). Irregularities of cleavage, such as prolonged or rapid cell cycles, may be associated with DNA repair activity, cellular rearrangement or failure to undergo cell cycle checkpoints (Regin et al., 2022).

As some significant timing differences have been reported with reference to specific outcome measures such as clinical pregnancy and chromosome complement, morphokinetic selection algorithms are being proposed to improve embryo selection and thereby, shorten the time to pregnancy (Meseguer et al.,

2011; Petersen et al., 2016; Pribenszky et al., 2017; Fishel et al., 2020). The potential of individual morphokinetic variables to predict clinical outcomes, has recently been assessed in two large analyses of over 30 000 embryos; the results show that periblastulation timings have more power to predict live birth than traditional TE or ICM morphology (Bamford et al., 2022; Campbell et al., 2022a). However, two recent randomized controlled trials (RCTs) found no improvement in ongoing pregnancy rate or cumulative live birth rate or live birth rate per transfer, when using TLT algorithmic selection (Ahlsröm et al., 2022; Kieslinger et al., 2023), corroborating the findings of the latest Cochrane review (Armstrong et al., 2019).

Although the studies are heterogeneous and drawing strong conclusions is difficult, TLT studies can help inform and optimize static assessment timing windows in the IVF laboratory. However, many laboratories do not have this technology, and the familiar, reliable daily descriptors remain practically applicable, although somewhat imprecise. Since the publication of the original Istanbul Consensus (2011), the convention of describing the timing of preimplantation development in terms of number of days (post insemination) has come to be viewed as simplistic, largely due to the facility to observe the developing embryo almost continuously, in minutes and hours, rather than days, using TLT imaging.

Consensus points

- Standardized timing of observations is critical for reliable comparison of results between different laboratories, culture conditions, patients, and other variables. This should be set relative to the time of insemination, and uniformly reported as hours post-insemination.
- There is an inherent variability in timing of all biological processes; the suggested observation times reflect those at which the associated developmental stages occur in most cases, whilst accepting there are confounding and influencing factors, including human subjectivity.
- Culture media and culture systems in general are recognized as having a significant impact on embryo morphokinetics; accordingly, their impact should be considered in comparative studies.
- Each laboratory is encouraged to develop and analyse its own datasets to determine relevant timings. Data generated by other laboratories may or may not be generally applicable.

2. Oocyte

Oocyte morphology may be assessed with the aim of predicting the developmental competence of the resulting embryo. In the relevant literature, several extra-cytoplasmic (cumulus oocyte complex (COC), ZP, perivitelline space (PVS), polar body (PB), shape, size) and intracytoplasmic (vacuoles, refractile bodies (RFs), aggregates of smooth endoplasmic reticulum clusters (sER-a), central granularity, colour) oocyte dysmorphic features are reported.

In this section, the predictive value of oocyte morphological characteristics/dysmorphism for embryo developmental potential is assessed (Table 2). Moreover, the possible use of oocytes that are immature at the time of oocyte retrieval following standard ovarian stimulation (so-called rescue in vitro maturation (rescue-IVM) is considered).

Oocyte morphological features relevant to oocyte scoring

The Istanbul Consensus (2011) described the optimal oocyte morphology as an oocyte with a spherical shape enclosed by a uniform ZP, with a uniform translucent cytoplasm free of inclusion, and a

Table 2. Overview of all evidence and recommendations for oocyte assessment.

Morphological feature	Atypical patterns		Summary of review findings		Considerations	Recommendation
	Fertilization rate	Blastocyst formation rate	Implantation rate	Live birth rate		
Cumulus oocyte complex (COC)	Compact COC Association with lower fertilization rate Very low ⊕○○○ 1 observational study (Rattanachaiyanont et al., 1999)	N/R	Association with lower pregnancy rate Very low ⊕○○○ 1 observational study (Dal Canto et al., 2012a)	N/R	Further studies are necessary before establishing the potential predictive value of this assessment on embryo competence	The presence of a dense COC and a very tight corona, if present in most of collected COCs from one patient, should be noted
	Presence of blood clots Associated with lower fertilization rate Very low ⊕○○○ 2 observational studies (Daya et al., 1990; Ebner et al., 2008a)	Associated with lower blastocyst formation Very low ⊕○○○ 1 observational study (Ebner et al., 2008a)	N/R	N/R		
Zona pellucida (ZP)	Dark/Thick ZP Contradictory results: No clear association with fertilization rate Very low ⊕○○○ 6 observational studies (De Sutter et al., 1996; Balaban et al., 1998; Esfandiari et al., 2006; Ten et al., 2007; Rienzi et al., 2008; Shi et al., 2014)	No clear association with blastocyst formation Very low ⊕○○○ 1 observational study (Balaban et al., 2008)	Contradictory results: No clear association with implantation rate Very low ⊕○○○ 3 observational studies (Esfandiari et al., 2006; Balaban et al., 1998; Pan and Zhang, 2020)	Association with lower live birth rate Very low ⊕○○○ 3 observational studies (Shi et al., 2014; Sauerbrun-Cutler et al., 2015; Sousa et al., 2015)	Evidence is insufficient to support any negative prognosis of zona pellucida characteristics/dysmorphisms on embryo developmental potential	Oocytes showing different ZP phenotypes are suitable for clinical use.
	Associated with lower fertilization rate Very low ⊕○○○ observational studies (Bertrand et al., 1995; Shi et al., 2014; Pan and Zhang, 2020)	N/A	Association with lower implantation rate Very low ⊕○○○ 3 observational studies (Shi et al., 2014; Sauerbrun-Cutler et al., 2015; Sousa et al., 2015)	N/R		
Perivitelline space (PVS)	Large PVS Association with lower fertilization rate Low ⊕⊕○○ 1 meta-analysis of 4 observational studies and 2 observational studies (Rienzi et al., 2008; Setti et al., 2011; Ashrafi et al., 2015)	No clear association with blastocyst formation rate Very low ⊕○○○ 1 observational study (Ferrarini Zanetti et al., 2018)	Association with lower implantation rate Very low ⊕○○○ 1 observational study (Kahraman et al., 2000; Ferrarini Zanetti et al., 2018)	N/R	Evidence is insufficient to support any negative prognosis of atypical PVS phenotype/size on embryo developmental potential	Oocytes showing different PVS phenotypes are suitable for clinical use.
	Granulated PVS No clear association with fertilization rate Very low ⊕○○○ A meta-analysis of 3 observational studies (Setti et al., 2011)	N/A	N/A	N/A		

(continued)

Table 2. Continued

Morphological feature	Summary of review findings				Recommendation		
	Atypical patterns	Fertilization rate	Blastocyst formation rate	Implantation rate			
Polar body (PB)	Fragmented PB	No association with fertilization rate Low ⊕⊕⊕⊕ 1 meta-analysis of 7 observational studies (Setti et al., 2011; Ashrafi et al., 2015)	Association with lower blastocyst formation Very Low ⊕⊕⊕⊕ 1 observational study (Zhou et al., 2016)	No clear association with implantation rate Very low ⊕⊕⊕⊕ 6 observational studies (Verlinsky et al., 2003; Giotti et al., 2004; De Santis et al., 2005; Chamayou et al., 2006; Ten et al., 2007; Zhou et al., 2016)	No clear association with ongoing/delivery rate Very low ⊕⊕⊕⊕ 1 observational study (Zhou et al., 2016)	Future quantitative studies are necessary to understand the potential negative impact of large polar bodies on embryo developmental potential	Oocytes showing fragmented or large PB are suitable for clinical use. Very large polar body could be associated with abnormal meiotic spindle configuration and deserve more attention
	Large PB	Association with lower fertilization rate Low ⊕⊕⊕⊕ 1 meta-analysis of 4 observational studies (Setti et al., 2011)	N/R	N/R	N/R		
Vacuolization	Presence of vacuoles	Association with lower fertilization rate Low ⊕⊕⊕⊕ 1 meta-analysis of 3 observational studies and 3 observational studies (Rienzi et al., 2008; de Cássia et al., 2010; Setti et al., 2011; Ashrafi et al., 2015)	Association with lower blastocyst formation rate Very low ⊕⊕⊕⊕ 2 observational studies (Ebner et al., 2005; Sousa et al., 2016)	N/R	N/R	Evidence was insufficient to support any negative prognosis on embryo developmental potential	Oocytes showing vacuoles are suitable for clinical use
Refractile bodies (RF)	Presence of RF	No clear association with fertilization rate Low ⊕⊕⊕⊕ 1 meta-analysis of 3 observational studies and 1 observational study (Setti et al., 2011; Takahashi et al., 2020)	No clear association with blastocyst formation Very Low ⊕⊕⊕⊕ 1 observational study (Takahashi et al., 2020)	No clear association with implantation rate Very low ⊕⊕⊕⊕ 2 observational studies (Balaban et al. 1998; Takahashi et al., 2020)	N/R	Evidence was insufficient to support any negative prognosis on this phenotype on further embryo developmental potential.	Oocytes showing refractile bodies are suitable for clinical use.
	Large RF (>5µm)	Association with lower fertilization rate Very low ⊕⊕⊕⊕ 1 observational study (Otsuki et al., 2007)	Association with lower blastocyst formation rate Very low ⊕⊕⊕⊕ 1 observational study (Otsuki et al., 2007)				

(continued)

Table 2. Continued

Morphological feature	Atypical patterns		Summary of review findings			Considerations		Recommendation
	Fertilization rate	Blastocyst formation rate	Implantation rate	Live birth rate				
Aggregates of Smooth Endoplasmic Reticulum Clusters (sER-a)	<p>Presence of sER-a</p> <p>No clear association with fertilization rate Low ⊕⊕⊕⊕ 10 observational studies (Otsuki et al., 2004; Ebner et al., 2008b; Sá et al., 2011; Hattori et al., 2014; Setti et al., 2016; Shaw-Jackson et al., 2016; Gurunath et al., 2019; Wang et al., 2021; Xu et al., 2022; Fang et al., 2022)</p>	<p>No clear association with blastocyst formation Low ⊕⊕⊕⊕ 9 observational studies (Ebner et al., 2008b; Sá et al., 2011; Hattori et al., 2014; Setti et al., 2016; Shaw-Jackson et al., 2016; Gurunath et al., 2019; Wang et al., 2021; Fang et al., 2022; Xu et al., 2022)</p>	N/R	N/R	N/R	N/R	<p>SER-a positive oocytes could be inseminated, based on a case-by-case evaluation</p>	
	<p>Central cytoplasmic granulation</p> <p>Association with lower fertilization rate Low ⊕⊕⊕⊕ 7 observational studies (Serhal et al., 1997; Balaban et al., 1998; Kahraman et al., 2000; Chamayou et al., 2006; Wilding et al., 2007; Rienzi et al., 2008; Yi et al., 2019)</p>	<p>Association with lower blastocyst formation rate Very low ⊕⊕⊕⊕ 1 observational study (Balaban et al., 2008)</p>	<p>Association with lower implantation rate Very low ⊕⊕⊕⊕ 1 observational study (Kahraman et al., 2000)</p>	N/R	N/R	<p>The difference was statistically insignificant, and the evidence was insufficient to support any negative prognosis of this phenotype on embryo developmental potential.</p>	<p>Oocytes showing cytoplasmic granularity are suitable for clinical use</p>	
Shape	<p>Ovoid oocyte</p> <p>No association with fertilization rate Very low ⊕⊕⊕⊕ 2 observational studies (Ebner et al., 2008c; Braga et al., 2013)</p>	<p>Association with lower blastocyst formation rate Very low ⊕⊕⊕⊕ 1 observational study (Ebner et al., 2008c)</p>	<p>No association with implantation rate Very low ⊕⊕⊕⊕ 5 observational studies (De Sutter et al., 1996; Balaban et al., 1998; Chamayou et al., 2006; Ten et al., 2007; Yakin et al., 2007)</p>	N/R	N/R	<p>Irregularly shaped oocytes are considered suitable for clinical use.</p>		
Colour	<p>Ooplasm darkness</p> <p>No association with fertilization rate Low ⊕⊕⊕⊕ 1 meta-analysis and 2 observational studies (Esfandiari et al., 2006; Setti et al., 2011; Shi et al., 2014)</p>	<p>Associated with lower blastocyst formation Very Low ⊕⊕⊕⊕ 1 observational study (Balaban et al., 2008)</p>	N/R	N/R	N/R	<p>Few studies investigated colour variation, often observed together with other anomalies.</p>	<p>Oocytes showing colour variation are suitable for clinical use.</p>	

(continued)

Table 2. Continued

Morphological feature	Summary of review findings				Considerations	Recommendation
	Atypical patterns	Fertilization rate	Blastocyst formation rate	Implantation rate		
Immaturity	Immature MI oocytes	Association with lower fertilization rate Low ⊕⊕○○ (De Vos et al., 1999; Balakier et al., 2004; Shu et al., 2007; Strassburger et al., 2010; Yang et al., 2021; Shani et al., 2023)	Association with lower blastocyst formation Very low ⊕○○○ 1 observational study (Yang et al., 2021)	N/R	Few live births obtained from rescue-IVM Very low ⊕○○○ 4 observational studies (Rubino et al., 2016; Escrich et al., 2018; Moon et al., 2023; Shani et al., 2023)	Due to their lower developmental potential, immature oocytes could be considered in case of poor prognosis individuals/couples and/or when alternatives are not available.
	Immature GV oocytes	No clear association with fertilization rate Very low ⊕○○○ 2 observational studies (Escrich et al., 2018; Shani et al., 2023)	No clear association with blastocyst formation Very Low ⊕○○○ 1 observational study (Escrich et al., 2018)			
Oocyte size	Oocyte with small ooplasm (<100 μm diameter)	Very low development potential Very low ⊕○○○ 1 observational study (Bassil et al., 2021)	N/R	N/R	N/R	Due to their lower developmental potential, very small oocytes could be considered only when alternatives are not available. It is recommended to exclude giant oocytes from all IVF/ICSI treatment programs due to their presumably possible tetraploid origin.
	Giant oocyte (>180 μm diameter)	Potential complications Very low ⊕○○○ 2 observational studies (Rosenbusch et al., 2002; Kitasaka et al., 2022)	N/R	N/R	N/R	

COC, cumulus oocyte complex; GV, Germinal vesicle; IVM, in vitro maturation; MI, Metaphase I; N/R, not reported; PB, polar body; PVS, perivitelline space; RF, refractile bodies; sER-a, aggregates of Smooth Endoplasmic Reticulum Clusters; ZP, zona pellucida.

Table colour code: Green: the oocyte can be clinically used; Yellow: the oocyte could be used with cautionary considerations. Red: the oocyte is not considered suitable for clinical use.

size-appropriate PB. Furthermore, it was noted that oocytes undergo both nuclear and cytoplasmic maturation, and that these processes are not equivalent, nor are they necessarily synchronous.

The survey results showed that 35% of respondents always apply the Istanbul Consensus (2011) recommendations to score oocytes, ranging from 22% for scoring the COC to 53% scoring the PB (Supplementary Data SII, Fig. 3B).

Cumulus oocyte complex

Most studies show an association between COC morphology and biological and clinical outcomes (Daya et al., 1990; Ng et al., 1999; Lin et al., 2003; La Sala et al., 2009; Dal Canto et al., 2012a). More specifically, the presence of a compact COC and a very tight corona has been found to be negatively associated with fertilization and pregnancy rates. On the other hand, no association was observed in one study between COC morphology and fertilization rate or embryo cleavage (Rattanachaiyanont et al., 1999). Further evidence indicates that the presence of blood clots trapped in the COC has a negative impact on outcomes even if removed during oocyte collection (Daya et al., 1990; Ebner et al., 2008a).

These data suggest that such COC characteristics, if present in most of collected COCs from one patient, should be noted, especially if conventional IVF (cIVF) is used for insemination. However, further studies are necessary before establishing the potential predictive value of this assessment for embryo competence.

Zona pellucida

Different ZP phenotypes (increased thickness, irregularities of the surface and increased density) have been reported. Some studies reported that oocytes with indented, thicker, dark and/or heterogeneous ZP had lower fertilization rate, embryo quality, embryo development, pregnancy, implantation, and live birth rates (Bertrand et al., 1995; Shi et al., 2014; Sauerbrun-Cutler et al., 2015; Sousa et al., 2015; Pan and Zhang, 2020; Yang et al., 2022). On the other hand, in several studies, ZP with diverse phenotypes showed no association with fertilization rates, embryo quality, implantation rates (De Sutter et al., 1996; Balaban et al., 1998; Esfandiari et al., 2006; Ten et al., 2007; Rienzi et al., 2008), embryo cryo-survival, or blastocyst and hatching rates (Balaban et al., 2008).

Only one study investigated the fertilization potential of oocytes without ZP (Ueno et al., 2014). Very rarely, two oocytes may share a single ZP. One live birth of dizygotic twins obtained from transfer of a pair of (zona-)conjoined blastocysts has been reported (Magdi, 2020). Moreover, two case reports described live births obtained from the transfer of embryos derived from insemination of (zona-)conjoined oocytes, one mature and the other immature (Fu et al., 2022a; Wang et al., 2022).

ZP birefringence, a refractive index derived from the polarization and propagation direction of light, has been utilized to predict the developmental potential of oocytes. Oocytes that exhibited high birefringence in the inner layer of the ZP were associated with higher implantation, pregnancy, and live birth rates compared to those with low birefringence in the inner layer of the ZP (Rama Raju et al., 2007; Montag et al., 2008; Madaschi et al., 2009). Moreover, the miscarriage rate was higher in embryos transferred from oocytes with low birefringence (Madaschi et al., 2009). On the contrary, another study indicated no significant differences between high and low birefringence in the inner layer of the ZP (Tabibnejad et al., 2018).

Evidence was insufficient to support any negative prognosis of ZP characteristics for embryo developmental potential. Oocytes showing different ZP phenotypes are therefore considered suitable for clinical use.

Perivitelline space

Contradictory reports are found in the literature assessing different PVS phenotypes and developmental competence (De Sutter et al., 1996; Balaban et al., 1998; Hassan-Ali et al., 1998; Farhi et al., 2002; Chamayou et al., 2006; Ten et al., 2007; Balaban et al., 2008; Rienzi et al., 2008; Ashrafi et al., 2015; Sauerbrun-Cutler et al., 2015; Ferrarini Zanetti et al., 2018; Weghofer et al., 2019). Three studies have focused in particular on large PVS and fertilization rate, finding a significant negative association (De Sutter et al., 1996; Xia, 1997; Ten et al., 2007; Rienzi et al., 2008; Setti et al., 2011; Ashrafi et al., 2015).

On the other hand, evidence was insufficient to support a negative prognosis for embryo developmental potential. Oocytes showing different PVS phenotypes are therefore considered suitable for clinical use.

Polar body

Large or fragmented PB are commonly reported. No significant association was found between PB fragmentation and fertilization. Although some studies showed an association between different PB phenotypes and early embryo development (Ebner et al., 2000; Chamayou et al., 2006; Fancsovits et al., 2006; Rienzi et al., 2008; Navarro et al., 2009; Zhou et al., 2016), no association with implantation or clinical pregnancy was reported (Verlinsky et al., 2003; Ciotti et al., 2004; De Santis et al., 2005; Ten et al., 2007; Liu et al., 2024).

Evidence was insufficient to support any negative prognosis of PB size and fragmentation on embryo developmental potential. Oocytes showing fragmented or large PB are therefore considered suitable for clinical use. However, a disproportionately large PB, although very rare, could be associated with abnormal meiotic spindle morphology or positioning, and deserves more attention.

Shape

Mature human oocytes generally have a spherical shape, nevertheless oocytes with ovoid shapes are reported. Overall, an ovoid shape does not appear to affect laboratory and clinical outcomes (De Sutter et al., 1996; Balaban et al., 1998; Chamayou et al., 2006; Ten et al., 2007; Yakin et al., 2007; Anagnostopoulou et al., 2022). In case of an ovoid oocyte that leads to planar arrangement of blastomeres at the 4-cell stage, further development up to blastocyst stage was found to be delayed (Ebner et al., 2008c).

Irregularly shaped oocytes are considered suitable for clinical use.

Oocyte size

Without consideration of the ZP thickness, small (<100 µm diameter) and large oocytes (≥125 µm diameter) have been reported to have very low developmental potential (Bassil et al., 2021).

Giant oocytes (e.g. >180 µm diameter) should be excluded from clinical use due to their possible tetraploid origin (Rosenbusch et al., 2002; Kitasaka et al., 2022). Presumably, these oocytes originally derive from the fusion of two primordial oocytes. This is suggestive of the presence of two diploid chromosome complements and an overall tetraploid oocyte constitution (Balakier et al., 2002; Rosenbusch et al., 2002; Munné et al., 2004). On the other hand, siblings of giant oocytes with normal diameter have been shown to have normal developmental potential (Machtinger et al., 2011; Lehner et al., 2015).

Vacuolization

Vacuoles are membrane-bound, translucent and fluid-filled cytoplasmic inclusions that appear at the end of oocyte maturation

(Otsuki et al., 2004; Sfountouris et al., 2018). Vacuoles can appear individually or in multiples (Fancsovits et al., 2011). Very large vacuoles (>25 µm) might distort the oocyte cytoskeletal structure, impairing sperm-oocyte signalling, sperm binding, meiotic resumption, and embryo development (Wallbutton and Kasraie, 2010; Dal Canto et al., 2017).

Different studies have shown that vacuolization is associated with lower fertilization rate, compromised embryo development, and lower blastulation and cryo-survival rates (Ebner et al., 2005; Balaban and Urman, 2006; Ebner et al., 2006; Ten et al., 2007; Balaban et al., 2008; Rienzi et al., 2008; de Cássia et al., 2010; Sousa et al., 2016). In particular, the association between the presence of vacuoles and lower fertilization was confirmed in a meta-analysis (Setti et al., 2011). However, in this analysis, evidence was insufficient to support any negative prognosis in relation to embryo developmental potential. Oocytes showing vacuoles are therefore considered for clinical use. In ICSI cases, however, care should be taken in avoiding injection of the sperm into a vacuole.

The so-called 'bull's-eye inclusion' is a distinct, smooth, spherical structure that encloses vesicles and is encircled by lipid droplets (Sousa et al., 2016). The impact of these structures on developmental potential remains unknown.

Refractile bodies

RFs consist of a mix of lipids and dense granular material. They exhibit a yellow autofluorescence typical of lipofuscin (Sathananthan, 1994). A small number of publications have investigated the predictive value of RF and embryo developmental potential (Alikani et al., 1995; De Sutter et al., 1996; Balaban et al., 1998; Ebner et al., 2000; Otsuki et al., 2004; Setti et al., 2011; Takahashi et al., 2020). A lower fertilization rate is associated with the presence of such phenotype, in particular if larger than 5 mm (Otsuki et al., 2007).

Although fertilization rate may be affected, the evidence was insufficient to support any negative prognosis of this phenotype for further embryo development. Oocytes showing RF are therefore considered suitable for clinical use.

Smooth endoplasmic reticulum clusters (sER-a)

From an ultrastructural standpoint, SER-a consist of tubular clusters surrounded by mitochondria that appear as more densely packed areas than the surrounding regions (Sá et al., 2011). SER-a have been described as potential biomarkers of oocyte quality. Numerous studies suggested lower fertilization (Sá et al., 2011; Massarotti et al., 2021), embryo quality (Ebner et al., 2008b; Sá et al., 2011; Braga et al., 2013; Massarotti et al., 2021; Wang et al., 2021) and pregnancy rates (Otsuki et al., 2004; Setti et al., 2016; Gurunath et al., 2019; Massarotti et al., 2021), and increased miscarriage rates (Otsuki et al., 2004; Ebner et al., 2008b; Braga et al., 2013). Moreover, in small studies, higher rates of perinatal complications and birth defects were reported as being associated with this dysmorphism (Otsuki et al., 2004; Ebner et al., 2008b; Akarsu et al., 2009; Sá et al., 2011; Mateizel et al., 2013; Sfountouris et al., 2018). Conversely, more recent studies and a meta-analysis reported no difference in fertilization rate, blastocyst formation rate, neonatal outcomes (Hattori et al., 2014; Shaw-Jackson et al., 2016; Itoi et al., 2017; Zhang et al., 2021; Fang et al., 2022) or euploidy rates (Xu et al., 2022; Mizobe et al., 2023; Wang et al., 2023); this body of evidence reinforces the recommendation, also supported by the Vienna Consensus (ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine, 2017) that clinical use of SER-a positive oocytes may be considered.

Granularity

Oocytes with central granulation have been associated with defective pronuclear morphology, reduced embryo quality (Ebner et al., 2008a; Rienzi et al., 2008), decreased cryo-survival rate, compromised embryo developmental competence (Balaban et al., 2008; Ebner et al., 2008a; Rienzi et al., 2008) increased aneuploidy rate (Wang et al., 2023), and lower ongoing pregnancy rate (Kahraman et al., 2000). In contrast, other studies and meta-analyses suggest that centrally localized cytoplasmic granulation might be a normal/typical oocyte morphological feature (Wilding et al., 2007; Setti et al., 2011; Yi et al., 2019). Currently, there are no studies investigating the potential of these oocytes to produce viable pregnancies. Available evidence is insufficient to support a negative prognostic value of this dysmorphism relevant to embryo developmental potential. Oocytes showing cytoplasmic granularity are therefore considered suitable for clinical use.

Colour

Limited studies have investigated translucency variation, often observed together with other anomalies. Some have suggested an association between ooplasm darkness and poorer embryo quality (Loutradis et al., 1999; Ten et al., 2007). However, this finding was not confirmed by other investigations (De Sutter et al., 1996; Balaban et al., 1998; Esfandiari et al., 2006; Balaban et al., 2008; Shi et al., 2014). The highly subjective nature of these observations as well as heterogeneity of the data preclude any conclusions. Oocytes showing variations in translucency are therefore considered suitable for clinical use.

Immaturity

After standard ovarian stimulation, approximately 15–20% of oocytes fail to extrude the first PB and reach the metaphase II (MII) stage, remaining at the metaphase I (MI) or germinal vesicle (GV) stages (ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine, 2017; ESHRE Clinic PI Working Group et al., 2021). Studies using polarized light microscopy have shown that some oocytes, despite showing a PB in the PVS, may still be immature, specifically being in the early Telophase I. At this stage, there is still a connection between the ooplasm and the forming PB, with the meiotic spindle of meiosis I positioned between the two separating cells (Rienzi et al., 2003; Petersen et al., 2009; Rienzi et al., 2012; Holubcová et al., 2019). Thus, only by observing the presence of the meiosis II spindle in the cytoplasm, oocyte meiotic maturity can be certainly assessed. More evidence is needed to clarify the importance of this assessment to predict embryo developmental fate (Rienzi et al., 2011; Tabibnejad et al., 2018; Halim et al., 2024).

Immature oocytes are usually not used for insemination and are discarded. However, in the case of poor prognosis patients and in patients with an unsynchronized follicle cohort, the use of immature oocytes that can mature after a period of *in vitro* culture (i.e. rescue-IVM oocytes) could contribute to the number of embryos obtained in each cycle, potentially increasing the overall chances of pregnancy (Shu et al., 2007). Several studies have shown that MI oocytes that mature within 2–6 h from denudation may be injected and may contribute to the number of available embryos (De Vos et al., 1999; Balakier et al., 2004; Shu et al., 2007). By contrast, overnight *in vitro* culture of MI and GV oocytes did not improve results. GV and MI oocytes that mature *in vitro* after 24 h have compromised results in terms of fertilization and blastocyst formation rates (Yang et al., 2021), most probably due to a higher risk of being chromosomally abnormal (Strassburger et al., 2010). TLT analysis has also confirmed that rescue-IVM oocytes

differ from their sibling MII oocytes in morphokinetic profile, showing a delay in the early stages of embryo development (Faramarzi et al., 2018; Margalit et al., 2019; Shani et al., 2023). However, the feasibility of the rescue-IVM approach is supported by some studies reporting a contribution to embryo yield, and few live births obtained using those embryos (Rubino et al., 2016; ESCRICH et al., 2018; Moon et al., 2023; Shani et al., 2023).

Due to their lower developmental potential, immature oocytes could be considered for clinical use in poor prognosis cases.

Oocyte morphology and morphokinetics

Some studies investigated a possible relationship between different cytoplasmic phenotypes and morphokinetics. Although not a standard procedure for oocyte assessment, ZP birefringence was shown in a recent study not to be correlated with embryo morphokinetics (Tabibnejad et al., 2018), while another study reported an early t5 in oocytes with high birefringence (Faramarzi et al., 2017). In the latter study, tPB2, t5 and t8 (time to extrusion of the second polar body (PBII) and development at the 5- and 8-cell stage, respectively), were associated with oocyte diameter, while PVS size showed no association with early development morphokinetics (Faramarzi et al., 2019). Finally, the incidence of failure of PBII extrusion and the incidence of mitotic cleavage failure in oocytes with SER-a were found to be significantly higher than that in oocytes without SER-a (Otsuki et al., 2018).

Overall, individual dysmorphic features may not be strongly associated with viability and development potential or clinical outcomes. However, it is possible that occurrence of two or more of these features together exerts a negative influence on outcomes (Alikani et al., 1995; Bartolacci et al., 2022).

Consensus points

- Giant oocytes should be excluded from clinical use.
- The use of small/large oocytes and IVM-rescued oocytes should be documented for prognostic and traceability purposes due to their apparently lower developmental potential.
- Embryos derived from MII oocytes free of large or multiple vacuoles, SER-a, disproportional shapes and very large first PBs should be prioritized for clinical use.
- Prenatal follow-up and the follow-up of babies born from oocytes with atypical phenotypes and rescue-IVM demands attention.

3. Zygote stage

TLT has revealed the complexity of morphokinetic changes occurring during normal (Payne et al., 1997; Mio and Maeda, 2008; Aguilar et al., 2014; Coticchio et al., 2018) and abnormal (Ezoe et al., 2022b; Wei et al., 2022) fertilization, leading to a more accurate and in-depth approach to fertilization assessment. Dynamic monitoring of this stage was previously inaccessible by static observation. Preimplantation genetic testing for aneuploidy (PGT-A) is also contributing to define the chromosomal constitution of zygotes with pronuclear abnormalities.

In this section, the optimal timing for zygote assessment and the significance of zygote characteristics for embryo developmental potential are reviewed.

Timing of zygote assessment

The Istanbul Consensus (2011) considered static fertilization assessment as 'straightforward, based on the observation of two polar bodies (PBs) and two pronuclei (PNs) at 17 ± 1 hpi'.

The survey results showed that 68% of respondents always apply the Istanbul Consensus (2011) recommendations to assess the zygote stage at 17 h ± 1 hpi (Supplementary Data SII, Fig. 3A).

Only one, albeit a very large, TLT study attempted to optimize the timing of PN observation (Barrie et al., 2021b). Monitoring more than 54 746 ICSI and 26 302 cIVF embryos, the number of 2PN zygotes was annotated at 30-min intervals, between 15 and 20 hpi. In both insemination groups, the interval with the highest proportion (>98%) of visible 2PN zygotes was 16.0–16.5 hpi. At later intervals, this rate progressively decreased, due to early PN breakdown (PNBD) in some zygotes.

Morphological features relevant to zygote assessment

The Istanbul Consensus (2011) described that the optimal fertilized oocyte is a spherical oocyte with two polar bodies, and two centrally located, juxtaposed pronuclei that are even sized, with distinct membranes (Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology, 2011). The pronuclei should have comparable numbers and size of nucleolar precursor bodies (NPBs) that are ideally clustered at the region of membrane juxtaposition of the two PN.

The survey results showed that 68% of the respondents always apply the Istanbul Consensus (2011) recommendation to score the pronuclear stage (Supplementary Data SII, Fig. 3B).

The predictive value of pronuclear stage features for embryo quality is discussed below (Table 3).

Zygote size

Oocyte and zygote size is usually reported as diameter, projected area or volume. Fertilized oocytes normally undergo progressive and moderate shrinkage during fertilization, also as a result of PBII extrusion (Liu et al., 2014). One study investigated this phenomenon, reporting a lack of association with live birth rate (Barberet et al., 2019). A more recent analysis suggested a negative correlation between zygote diameter/cytoplasmic volume observed at 17 hpi and blastocyst quality (Kljajic et al., 2023). Collectively, this evidence is insufficient and inconclusive on the hypothesis that zygote size can be a predictive parameter for embryo developmental potential.

Pronuclei (PN)

Position. Using TLT, two studies investigated PN position as a developmental biomarker. Although rarely observed, off-centre position annotated shortly before PNBD was associated with abnormal division, namely trichotomous cleavage (Coticchio et al., 2018). Off-centre position of PNs at the time of juxtaposition (8–9 hpi) was found to be associated with a two-fold decrease in live birth rate (Barberet et al., 2019), also after multivariate analysis. Notably, the feature observed in the latter study cannot be detected by single static observation at 16–17 hpi.

Juxtaposition. In one TLT report, lack of PN juxtaposition throughout fertilization was observed in 1–2% of zygotes. In this phenotype, cleavage, morula, and blastocyst formation rates were negatively affected (Ezoe et al., 2022a).

Size. PNs increase in size progressively as soon as they form, reaching their final size shortly before PNBD (Otsuki et al., 2017; Orevich et al., 2022). TLT investigation confirmed that the paternal PN is normally larger than its female counterpart (Barberet et al., 2019; Ezoe et al., 2022b; Orevich et al., 2022). Size difference between the two PN tends to progressively decrease as fertilization unfolds. If assessed in the 16–18 hpi interval or immediately before PNBD, this difference was smaller in zygotes that resulted in live births (Otsuki et al., 2017; Otsuki et al., 2019). In addition, results from TLT are not conclusive on the value of PN size as an

Table 3. Overview of all evidence and recommendations for zygote assessment.

Morphological feature	Summary of review findings and level of evidence per outcome						Recommendation
	Atypical patterns	Abnormal cleavage rate	Cleavage rate	Blastocyst formation rate	Implantation rate	Live birth rate	
Zygote size	Diameter <113 µm	N/R	N/R	Association with lower blastocyst formation rate Very low ⊕○○○ 1 observational study (Kijajic et al., 2023)	N/R	No clear association with live birth rate Very low ⊕○○○ 1 observational study (Barberet et al., 2019)	Numerous zygotic attributes—zygote size, PN size, PN position, NPB patterning—might be associated with embryo quality and clinical outcome. Lack of PN juxtaposition is very rare, but strongly associated with poor blastocyst development.
PN position	Off-centre position	Association with higher abnormal cleavage rate Very low ⊕○○○ 1 observational study (Coticchio et al., 2018)	N/R	N/R	N/R	N/R	Abnormalities in PN position, juxtaposition and size are very rare and difficult, or impossible, to monitor by static observation. The evidence is insufficient for the application of the studied features as biomarkers.
	Off-centre juxtaposition	N/R	N/R	N/R	N/R	Association with lower live birth rate Very low ⊕○○○ 1 observational study (Barberet et al., 2019)	
	Lack of PN juxtaposition	Association with higher abnormal cleavage rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2022a)	Association with lower cleavage rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2022a)	Association with lower blastocyst formation rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2022a)	N/R	No clear association with live birth rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2022a)	
	Interpronuclear Difference in male and female PN areas	N/R	N/R	N/R	N/R	Association with lower live birth rate Very low ⊕○○○ 2 observational studies (Otsuki et al., 2017, 2019)	
Nucleolar precursor bodies	NPB patterns (Z1-Z4)	N/R	N/R	Association with higher blastocyst formation rate Very low ⊕○○○ 1 observational study (Cavazza et al., 2021)	No clear association with implantation rate Very low ⊕○○○ 1 observational study (Aguilar et al., 2014)	No clear association with live birth rate Very low ⊕○○○ 2 observational studies (Azzarello et al., 2012; Barberet et al., 2019)	The intrinsic morphological mutability during short time periods (NPB patterning) is not amenable to static observation
	Migration speed	N/R	N/R	N/R	N/R	Association with higher live birth rate Very low ⊕○○○ 2 observational studies (Inoue et al., 2017, 2019)	

(continued)

Table 3. Continued

Morphological feature	Summary of review findings and level of evidence per outcome					Recommendation	
	Atypical patterns	Abnormal cleavage rate	Cleavage rate	Blastocyst formation rate	Implantation rate		Live birth rate
Cytoplasmic halo	Absence of cytoplasmic halo	Association with higher abnormal cleavage rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2020)	Association with lower cleavage rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2020)	Association with lower blastocyst formation rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2020)	N/R	2021; Inoue et al., 2023) No clear association with live birth rate Very low ⊕○○○ 2 observational study (Barberet et al., 2019; Ezoe et al., 2023)	The absence of the cytoplasmic halo may be de-select, embryos in day 3 embryo transfers.
Number of PNs	OPN	N/R	Association with lower cleavage rate Very low ⊕○○○ 1 observational study (Fu et al., 2022b)	No clear association with blastocyst formation rate Very low ⊕○○○ 1 observational study (Fu et al., 2022b)	No clear association with implantation rate Low ⊕○○○ 3 observational studies (Liu et al., 2016; 2020; Fu et al., 2022b)	No clear association with live birth rate Low ⊕○○○ 4 observational studies (Liu et al., 2016; Destouni et al., 2018; Li et al., 2021; Fu et al., 2022b)	The term “OPN” should not be used, if based on static observation. “Not observed 2PN” or “not reported 2PN” may be alternative definitions of normal zygotes undergoing early PNBD and, for such a reason, not detected by static observation.
	1PN	Association with higher abnormal cleavage rate Very low ⊕○○○ 1 observational study (Ezoe et al., 2020)	No clear association with cleavage rate Low ⊕○○○ 2 observational studies (Capalbo et al., 2017; Fu et al., 2022b)	Association with lower blastocyst formation rate Low ⊕○○○ 3 observational studies (Itoi et al., 2015; Capalbo et al., 2017; Ezoe et al., 2022b)	No clear association with implantation rate Low ⊕○○○ 6 observational studies (Itoi et al., 2015; Hondo et al., 2019; Si et al., 2019; Si et al., 2020; Li et al., 2021; Fu et al., 2022b)	No clear association with live birth rate Low ⊕○○○ 6 observational studies (Itoi et al., 2015; Hondo et al., 2019; Si et al., 2020; Li et al., 2021; Fu et al., 2022b)	The evidence suggests a possible cautious clinical use of 1PN zygotes, combining blastocyst culture and -if available—PGT-A technology appropriate for biparental diploidy assessment
	2.1PN	N/R	Association with lower cleavage rate Very low ⊕○○○ 1 observational study (Capalbo et al., 2017)	Association with lower blastocyst formation rate Very low ⊕○○○ 1 observational study (Capalbo et al., 2017)	N/R	N/R	The clinical use of 2PN zygotes with one small micropronucleus (2.1 PN) may be considered, especially if associated with PGT-A technology appropriate for biparental diploidy assessment
	3PN	N/R	N/R	N/R	N/R	N/R	The clinical use of 3PN zygotes is not recommended, while pre-clinical studies should be encouraged

NBP, nucleolar precursor bodies; N/R, not reported; PGT-A, preimplantation genetic testing for aneuploidy; PN, pronucleus.

Table colour code: Green: the embryo can be clinically used; Yellow: the embryo could be used with cautionary considerations. Red: the embryo is considered not suitable for clinical use.

independent parameter associated with outcome. Collectively, these studies suggest that abnormalities in PN position, juxtaposition and size are very rare and difficult, or near impossible, to monitor by static observation.

Nucleolar precursor bodies

NPBs are intra-pronuclear aggregates of fibrillar material of largely unknown composition. Once condensed from amorphous material, they increase in size and finally cluster in the region of PN juxtaposition. NPB condensation and clustering reflects the distribution of zygotic chromatin (Cavazza et al., 2021). Chromatin remodelling may be a pre-requisite for optimal chromosome-spindle microtubules interaction and, ultimately, chromosome congression. TLT evidence on NPBs is not consistent. Studies focusing on implantation and live birth did not indicate a predictive value of NPB patterning (Azzarello et al., 2012; Aguilar et al., 2014; Barberet et al., 2019), unless NPB speed was assessed with complex computational methodology (Inoue et al., 2021; Inoue et al., 2023). Another recent investigation (Cavazza et al., 2021) suggested a positive association between NPB clustering in both PN in the regions of juxtaposition and higher competence for blastocyst development, confirming previous data from static observation (Tesarik and Greco, 1999). Such contradictions are expected. In fact, NPB clustering is a continuum that follows different kinetics in male and female PN (Mio and Maeda, 2008; Coticchio et al., 2018) and, once achieved, can even be lost due to active NPB dispersal in the few hours preceding PNBD (Cavazza et al., 2021). This complicates the use of NPB patterning as biomarker for embryo quality.

Cytoplasmic halo

The cytoplasmic halo is described as a cortical domain of the zygote denoted by reduced cytoplasmic granularity. Visible in most zygotes (82–98%), it can be symmetrically or asymmetrically positioned (Ebner et al., 2003). Usually, the halo forms 2–4 h after PN appearance and disappears ~1 h before PNBD (Coticchio et al., 2018; Ezoe et al., 2020). Its formation is probably due to centripetal displacement of mitochondria and other organelles towards the area surrounding the PNs (Squirrell et al., 2003). One TLT study including 1009 zygotes focused specifically on this feature and found that absence of the halo was strongly associated with abnormal cleavage and embryo attrition at cleavage and morula stages. However, in single vitrified-warmed embryo transfers, halo-positive and halo-negative blastocysts produced comparable clinical outcomes (Ezoe et al., 2020). In the same study, halo position (symmetric or asymmetric) was not correlated with laboratory or clinical outcomes. Another TLT analysis confirmed that live birth rate is unaffected in transfers of halo-negative embryos (Barberet et al., 2019). This evidence disputes the significance of the halo, especially if embryo culture is extended to the blastocyst stage.

Nulli- mono- and tri- pronuclear zygotes

A designation of normal fertilization typically relies on observation of two PN. However, in the past several years zygotes with other pronuclear patterns, discernible at the time of static fertilization assessment, have been considered for clinical use: no visible PN (0PN), one PN (1PN) or three PN (3PN). A fourth rarer profile showing 2PN with one (or more) extra micro-pronucleus, referred to as 2.1PN, has been also occasionally reported.

0PN. Overall morphokinetic evidence does not confirm that embryo development can occur in the absence of formation of at least one PN. Rather, in all likelihood, '0PN zygotes' progressing to the first mitosis are 2PN or, rarely, 1PN/multi-PN zygotes

undergoing PNBD before static fertilization assessment can detect PN presence (Barrie et al., 2021b). Therefore, it is not surprising that studies on '0PN zygotes' (all based on static fertilization assessment, here only a few cited) reported rates of development, euploidy, implantation and live births comparable with or higher than those of 2PN zygotes (Liu et al., 2016; Destouni et al., 2018; Hondo et al., 2019; Paz et al., 2020; Fu et al., 2021; Li et al., 2021; Kemper et al., 2023). In fact, in general, embryos displaying faster morphokinetics as early as the fertilization stage are also developmentally more competent (Coticchio et al., 2023).

1PN. The Vienna Consensus recommended that 1PN rate should not exceed 3% and 5% in cIVF and ICSI cycles, respectively (ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine, 2017). In unselected 1PN-derived ICSI embryos, all morphokinetic times and developmental rates are significantly affected (Ezoe et al., 2022b). However, in IVF/ICSI 1PN zygotes showing a relatively larger PN size (defined by projected area or diameter cut-offs of $\geq 710 \mu\text{m}^2$ and $\geq 31 \mu\text{m}$, respectively), cleavage and blastocyst formation rates are comparable with those of 2PN fertilization (Araki et al., 2018; Kai et al., 2018). It is plausible that a larger size of the single PN reflects a higher, possibly diploid, DNA content. Indeed, in ~50% of cases of monopronuclear fertilization following IVF, the presence of both maternal and paternal DNA inside the single PN was documented (Cohen et al., 1995; Kai et al., 2015). The genesis of biparental diploid 1PN zygotes may differ in cIVF and ICSI fertilization. A recent TLT investigation suggests a possible modality of formation of biparental 1PN zygotes in cIVF: if, at the very beginning of fertilization, the fertilizing sperm penetrates the oocyte near (within a radius of $18 \mu\text{m}$) the presumed position of the maternal chromosomes, as suggested by the PBII localization, the paternal and maternal chromatin may be recruited together in the formation of a single PN (Wei et al., 2022). Consistent with this, several studies reported that 1PN blastocysts screened by PGT-A were diploid/euploid in significant proportions (40–50% of tested samples), in some cases, similar to those of 2PN controls (Bradley et al., 2017; Capalbo et al., 2017; Destouni et al., 2018; Xie et al., 2018; Zhao et al., 2022). In addition, while such studies involved ICSI as part of the PGT-A procedure, live births from 1PN zygotes have also been obtained in cIVF cases (Li et al., 2020). Documented use of 1PN zygotes for clinical purposes have been numerous (here only a few are reported). Overall, following blastocyst culture adopted to select more developmentally competent embryos, rates of implantation, pregnancy, and live birth approached those derived from 2PN zygotes (Itoi et al., 2015; Hondo et al., 2019; Si et al., 2019; Li et al., 2020; Li et al., 2021; Fu et al., 2022b; Kemper et al., 2023).

3PN. According to the recommendations of the Vienna Consensus, polypronuclear (including 3PN) fertilization should be <6% (ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine, 2017). Morphokinetics and blastocyst development of 3PN zygotes is less affected compared with 1PN fertilization (Ezoe et al., 2022c). The origin of 3PN zygotes may be digynic or di/polyandric, also depending on the type of insemination technique. Reports on PGT-A analysis and clinical use of 3PN zygotes are very rare. In a study based on 30 3PN blastocysts the rate of diploidy/euploidy was 33% (Mutia et al., 2019). In a case report, an apparently healthy live birth was achieved from the transfer of one euploid 3PN blastocyst (Yalçınkaya et al., 2016). A recent report described a healthy live birth and normal postnatal development up to 4 years from the transfer of a 4PN zygote (Bredbacka et al., 2023). However, in presumptive 3PN/4PN zygotes the origin of the third/fourth PN

(whether true extra PN or 'larger than usual' micropronucleus) remains a matter of ambiguity.

Micropronuclei. At the time of PN assessment, one or more small extra PNs may be rarely observed. They may originate from assembly of one extra small nuclear compartment around one or more chromosomes of a diploid zygote (Currie et al., 2022). Specific TLT investigations are lacking. One study based on static observation and PGT-A monitored >3500 zygotes, among which only <1% (n = 27) were 2PN showing one small extra PN (referred to as 2.1PN zygotes) (Capalbo et al., 2017). Although these zygotes show reduced first cleavage rate (74%), they can develop into biparental diploid blastocysts and produce apparently normal live births.

Consensus points

- Evidence reveals considerable plasticity of human fertilization and provides the basis for updated recommendations relevant to static fertilization assessment.
- **Timing of observation.** For static observations, assessment of PN number should be carried out at 16–17 hpi in both cIVF and ICSI cases, to minimize the probability that zygotes undergoing relatively early PNBD are incorrectly classified as unfertilized oocytes. Checking for syngamy (disappearance of PN) by static observation, mentioned in the Istanbul Consensus (2011), is not recommended since timing of PNBD cannot be precisely determined.
- **Morphological features.** Numerous zygotic attributes, including zygote size, PN size, PN position and NPB patterning, may be associated with embryo quality and clinical outcome. However, their use as biomarkers is hindered by at least two factors: (i) insufficient evidence (e.g. PN size), and (ii) intrinsic morphological mutability during short time periods (NPB patterning) not amenable to static observation. Lack of PN juxtaposition is very rare, but strongly associated with poor blastocyst development. The absence of the cytoplasmic halo affects blastocyst formation, but not implantation rate after blastocyst transfer. Therefore, the absence of the halo may be used to rank, but not de-select, embryos in Day-3 embryo transfers.
- **PN number.** By static observation, pronuclei may not be seen at fertilization check, and yet normal embryo development can occur. This may be explained by TLT data, which show that a significant proportion of 2PN zygotes undergo PNBD at earlier times than the fertilization check interval recommended by the original Istanbul Consensus (2011). In such cases, the presence of the PBII should accompany 2PN fertilization and therefore be used as a scoring criterion. While these zygotes may be categorized as OPN, if cultured, they may produce normal laboratory and clinical outcomes. Therefore, the term unfertilized or 'OPN' should not be used in these cases. Instead, 'PN not observed' may be a more suitable alternative for zygotes undergoing normal development without confirmation of fertilization.

Preliminary PGT-A data suggest that a significant proportion of 1PN and, some 3PN zygotes may be biparental diploid. In addition, a growing number of studies have reported normal live births from 1PN zygotes derived from both ICSI and IVF cycles. Collectively, this evidence supports cautious clinical use of 1PN zygotes, combining blastocyst culture and, if available, PGT-A technology appropriate for biparental diploidy assessment. The clinical use of 3PN zygotes is not recommended based on current evidence. 2PN zygotes with one extra micropronucleus (2.1PN) are relatively rare. However, they also may have a diploid

genotype and lead to apparently normal live births. Their clinical use may be considered, especially if associated with PGT-A technology. In general, the possible clinical use of 1PN and 2.1PN zygotes should be discussed with the clinical team and the patient, and governed by an internally approved policy.

4. Cleavage stage

Assessment of embryos at predefined times on Days 1, 2 and 3 has shown number of cells, fragmentation grade, blastomere size, and multinucleation to correlate with pregnancy and live birth outcomes (Lundin and Ahlström, 2015).

The survey results indicate that the vast majority of clinics (95%) still perform early-stage embryo evaluations. However, the traditionally static 'snapshot' assessments once or twice per day implies that no information regarding the development between these time points is obtained. Therefore, significant events such as abnormal cell divisions may be missed. Also, it has been shown that the morphology of an embryo may change in a couple of hours, for a better or a worse score (Montag et al., 2011), one reason being the dynamic occurrence and reabsorption of fragments during the cleavage process (Hardarson et al., 2001).

This section discusses morphological and morphokinetic attributes assessed at the early embryo cleavage stages and their potential impact on success rates for an embryo transferred or cryopreserved on Day 2 or Day 3 post fertilization (Table 4). It is important to consider that the same attributes may not be relevant or may have a different impact if the embryo survives extended culture and is transferred, fresh or after cryopreservation, at the blastocyst stage.

Timing of cleavage-stage embryo assessment

The Istanbul Consensus (2011) recommended static observation performed at 44 ± 1 hpi for Day-2 embryos and 68 ± 1 hpi for Day-3 embryos. The survey results showed that 41% and 63% of the respondents always assessed embryos on Day 2 or Day 3, respectively, applying these recommendations (Supplementary Data SII, Fig. 3A).

Assessment by TLT permits more detailed analysis of the traditional morphological parameters over time, as well as the incidence of abnormal cleavages. Several early, retrospective, TLT studies found that morphokinetic variables such as the timing of the first cell division, as well as the lengths of cell cycles, correlated with further embryonic development and subsequent pregnancy outcomes (Meseguer et al., 2011; Dal Canto et al., 2012b; Herrero et al., 2013). However, recent RCTs and meta-analyses have not found improvement in live birth rates following embryo selection using TLT algorithms (Armstrong et al., 2019; Ahlström et al., 2022; Kieslinger et al., 2023).

More recent TLT studies have shown timings with slight deviations from those reported in the Istanbul Consensus (2011), the differences becoming more pronounced and varied from the 4-cell stage onwards (Table 1).

Timing of first cleavage

The single most important indicator of embryo viability is cellular division. The occurrence of early cleavage, i.e. the first cell division occurring before 25–27 hpi, has been shown to correlate positively with embryo quality on Day 2 and Day 3, blastocyst formation rate (Herrero et al., 2013; de los Santos et al., 2014; Milewski et al., 2015), and implantation and live birth rates after transfer on Day 2 or 3 (Lundin et al., 2001; Salumets et al., 2003). This is also more recently supported by TLT studies (Coticchio et al., 2018; Sayed et al., 2020). In addition, TLT has shown that the time from disappearance of pronuclei or pronuclei fading

Table 4. Overview of all evidence and recommendations for cleavage stage embryo assessment.

Morphological feature	Atypical pattern		Summary of review findings		Considerations	Recommendation
	Embryo quality and development potential	Ploidy	Implantation rate	Live birth rate		
First cleavage	<p>Association with higher embryo quality and blastocysts formation rate Very low ⊕○○○ (Herrero <i>et al.</i>, 2013; de los Santos <i>et al.</i>, 2014; Milewski <i>et al.</i>, 2015)</p>	<p>Association with higher aneuploidy rate Very low ⊕○○○ (Vera-Rodriguez <i>et al.</i>, 2015)</p>	<p>Contradictory results: No association with implantation rate Low ⊕⊕○○ (5 observational studies (Lundin <i>et al.</i>, 2001; Salumets <i>et al.</i>, 2003; Ahlström <i>et al.</i>, 2016; Coticchio <i>et al.</i>, 2018; Sayed <i>et al.</i>, 2020) Association with higher implantation rates Low ⊕⊕○○ (1 RCT and 3 observational studies (Thurin <i>et al.</i>, 2005; Sundström and Saldeen, 2008; De los Santos <i>et al.</i>, 2014; Yang <i>et al.</i>, 2015))</p>	<p>Contradictory results: No association with live birth rate Low ⊕⊕○○ (1 RCT and 3 observational studies (Thurin <i>et al.</i>, 2005; Salumets <i>et al.</i>, 2003; Ahlström and Saldeen, 2008; De los Santos <i>et al.</i>, 2014; Yang <i>et al.</i>, 2015) Association with higher live birth rates Low ⊕⊕○○ (5 observational studies (Lundin <i>et al.</i>, 2001; Salumets <i>et al.</i>, 2003; Ahlstrom <i>et al.</i>, 2016; Coticchio <i>et al.</i>, 2018; Sayed <i>et al.</i>, 2020))</p>	<p>Assessment of early cleavage embryos may add information regarding other features such as binucleation/multinucleation and cell size. An important aspect to consider is the difference between zygotes originating from ICSI or cIVF.</p>	<p>The importance of scoring early cleavage for prediction of success rates has not been conclusively established.</p>
Abnormal early cleavage (direct cleavage, reverse cleavage, irregular chaotic division)	<p>N/R</p>	<p>Association with higher aneuploidy rate Low ⊕⊕○○ (3 observational studies (Arroyo <i>et al.</i>, 2015; Desai <i>et al.</i>, 2018))</p>	<p>Association with lower implantation rate Low ⊕⊕○○ (4 observational studies (Meseguer <i>et al.</i>, 2011; Petersen <i>et al.</i>, 2016; Zhan <i>et al.</i>, 2016; Liu <i>et al.</i>, 2020))</p>	<p>N/R</p>	<p>Assessment of early cleavage by TLT can be used to select against abnormal cleavage patterns such as direct cleavage, reverse cleavage, and irregular chaotic division.</p>	<p>Assessment of early cleavage by TLT can be used to select against abnormal cleavage patterns such as direct cleavage, reverse cleavage, and irregular chaotic division.</p>
Cell numbers	<p>Association with embryo scoring Very low ⊕○○○ (3 observational studies (Alikani, 2003; Machtinger and Racowsky, 2013; Yu <i>et al.</i>, 2018))</p>	<p>Correlation with chromosomal status Low ⊕⊕○○ (3 observational studies (Almeida and Bolton, 1996; Magli <i>et al.</i>, 2007; Kroener <i>et al.</i>, 2015))</p>	<p>Correlation with implantation rates Low ⊕⊕○○ (4 observational studies (Giorgetti <i>et al.</i>, 1995; Alikani <i>et al.</i>, 2000; Van Royen <i>et al.</i>, 2001; Rhenman <i>et al.</i>, 2015))</p>	<p>Correlation with live birth rates Low ⊕⊕○○ (5 observational studies (Giorgetti <i>et al.</i>, 1995; Racowsky <i>et al.</i>, 2011; Rhenman <i>et al.</i>, 2015; Awadalla <i>et al.</i>, 2022b; Tian <i>et al.</i>, 2022))</p>	<p>The current expected observation for embryo development is 4 cells on Days 2 and 8 cells on Day 3.</p>	<p>The current expected observation for embryo development is 4 cells on Days 2 and 8 cells on Day 3.</p>
Fragmentation	<p>Association with embryo quality and development potential Very low ⊕○○○ (2 observational studies (Alikani <i>et al.</i>, 2000; Ebner <i>et al.</i>, 2001))</p>	<p>Association with lower euploidy rate Very low ⊕○○○ (3 observational studies (Munné <i>et al.</i>, 1995; Ziebe <i>et al.</i>, 2003; Chavez <i>et al.</i>, 2012))</p>	<p>Association with lower implantation rate Very low ⊕○○○ (4 observational studies (Alikani <i>et al.</i>, 1999; Ebner <i>et al.</i>, 2001; Van Royen 2001; Racowsky <i>et al.</i>, 2011))</p>	<p>Association with lower live birth rates Low ⊕⊕○○ (3 observational studies (Rhenman <i>et al.</i>, 2015; Ahlstrom <i>et al.</i>, 2016; Awadalla <i>et al.</i>, 2022b))</p>	<p>The percent values are based on the cell equivalents, so for a 4-cell embryo, 25% fragmentation would equate to one blastomere in volume.</p>	<p>The relative degrees of fragmentation were defined as: no or minimal (<10%), mild (≤25%), or severe (>25%).</p>

(continued)

Table 4. Continued

Morphological feature	Summary of review findings			Considerations		Recommendation
	Atypical pattern	Embryo quality and development potential	Ploidy	Implantation rate	Live birth rate	
Cell size	Uneven cellular cleavage	N/R	Correlation with chromosomal errors Low ⊕⊕○○ 2 observational studies (Hardarson et al., 2001; Shenoy et al., 2021)	Association with lower implantation rate Very low ⊕○○○ 1 observational study (Múgica et al., 2008)	N/R	It is important to consider that the relative cell sizes must be 'cell stage appropriate', i.e. assessed in relation to the number of cycles that they have gone through For embryos at the 2-, 4-, and 8-cell stages, blastomeres should be evenly sized. For all other cell stages, one would expect a cell stage appropriate size difference as the cleavage phase has not been completed.
Multinucleation	Multiple nuclei	Negative correlation with time of development Low ⊕⊕○○ 5 observational studies (Ergin et al., 2014; Goodman et al., 2016; Balakier et al., 2017; Desch et al., 2017; Sayed et al., 2022)	No association with aneuploidy rates Very low ⊕○○○ 1 observational study (Desai et al., 2018)	Association with lower implantation rate Low ⊕⊕○○ 4 observational studies (Ergin et al., 2014; Goodman et al., 2016; Desch et al., 2017; Sayed et al., 2022)	Association with lower live births rate Low ⊕⊕○○ 4 observational studies (Ergin et al., 2014; Goodman et al., 2016; Desch et al., 2017; Sayed et al., 2022)	Multinucleation assessment on Day 3 would be complicated by the much smaller cell size, and therefore would be less reliable. If available, multinucleation should be scored using TL.T. Laboratories should record the incidence and discriminate between binucleation, multinucleation and micronucleation in each embryo, and ideally, the nucleation status of each blastomere in each embryo.
	Binucleation and/or micronucleation	Association with higher blastocyst formation rate Very low ⊕○○○ 1 observational study (Talbot et al., 2022)	N/R	Association with higher implantation rate Very low ⊕○○○ 2 observational studies (Aguilar et al., 2016; Talbot et al., 2022)	N/R	Binucleation on Day 2, at the 4-cell stage, may not be necessarily a negative sign, but more evidence is needed.
Other morphological features	Spatial disorganization	No clear association with embryo development Very low ⊕○○○ 2 observational studies (Ebner et al., 2012, 2017)	N/R	N/R	N/R	Embryos with apparent spatial disorganization should not be considered abnormal. There is no significant body of evidence to support a clear biological effect of cytoplasmic granularity, membrane appearance and the presence of vacuoles, these features on implantation potential.
	Cytoplasmic granularity, membrane appearance, vacuoles	Negative correlation with Day 3 development (atypical early compaction) Low ⊕⊕○○ 3 observational studies (Skiadas et al., 2006; Le Cruguel et al., 2013; Aslan Öztürk et al., 2022)	N/R	N/R	N/R	More research is required to identify which, if any, of these features are correlated with (or indicative of) implantation potential

ciVF, Conventional In Vitro Fertilization; N/R, not reported.

Table colour code: Green: the embryo can be clinically used; Yellow: the embryo could be used with cautionary considerations. Red: the embryo is not considered suitable for clinical use.

(PNf) to the start of the first cytokinesis was significantly related to ploidy (Vera-Rodriguez et al., 2015). A retrospective analysis of Day-2 single embryo transfers of ICSI embryos ($n=207$), including both traditional morphology variables as well as morphokinetic variables and patient characteristics, showed early cleavage, measured as more than one cell at 25–27 hpi, to be a significant predictor of live birth (OR 4.84, CI 2.14–10.96, $P=0.0002$) (Ahlstrom et al., 2016). In addition, it was found that each increase in grade of fragmentation (to 5–10%, 11–20%, 21–50%, 51–100%) significantly decreased the probability for live birth (OR 0.46, CI 0.25–0.84, $P=0.012$).

The same study also found that, for Day-2 transfers, early cleavage and fragmentation grade were better predictors of live birth outcome when compared with morphokinetic variables, and that no morphokinetic variables up to Day 2 improved prediction of live birth further (Ahlstrom et al., 2016). However, other studies have not found any correlation between early cleavage and implantation or live birth (Thurin et al., 2005; Sundström and Saldeen, 2008; de los Santos et al., 2014; Yang et al., 2015), and the data on potential importance of scoring early cleavage are currently inconclusive.

Still, the assessment of early cleavage in a TLT system can be used to select against abnormal early cleavages such as direct cleavage, reverse cleavage and irregular chaotic division, which have been shown to be associated with lower blastocyst formation rates, implantation and live birth rates (Meseguer et al., 2011; Petersen et al., 2016; Zhan et al., 2016; Liu et al., 2020) as well as with aneuploidy (Arroyo et al., 2015; Yan et al., 2015; Desai et al., 2018) and multinucleation (Zhan et al., 2016). In a study by Barrie et al., the prevalence of these abnormal cleavages was found to be 11.4% per cleaved embryo (Barrie et al., 2017b).

At present, the use of early cleavage/early syngamy in scoring regimens varies greatly between laboratories. An important aspect to consider is the difference between zygotes originating from ICSI and cIVF, as discussed in Section 1 (Expected timeline of embryo development and morphology) and Section 3 (Zygote stage assessment) of this paper.

Number of cells on Day 2 and Day 3

The number of blastomeres at a specific time signifies the developmental rate of the embryo and is considered the most important parameter for embryo scoring (Machtinger and Racowsky, 2013; Yu et al., 2018). Many earlier studies already showed the number of cells at Day 2 or Day 3 to be highly predictive of laboratory and clinical outcomes (Giorgetti et al., 1995; Alikani et al., 2000; Holte et al., 2007; Racowsky et al., 2011).

The Istanbul Consensus (2011) defined an optimal Day-2 embryo (44 ± 1 hpi) as an embryo with 4 equally sized mononucleated blastomeres in a three-dimensional tetrahedral arrangement, with <10% fragmentation, and a Day-3 embryo (68 ± 1 hpi) with 8 equally sized mononucleated blastomeres in a three-dimensional tetrahedral arrangement, with <10% fragmentation (Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology, 2011). The survey results showed that 68% of the respondents apply these Istanbul Consensus (2011) recommendations to score Day-2 and Day-3 embryos (Supplementary Data SII, Fig. 3B).

There seems to exist an 'optimal' development speed and many publications throughout the years have reported that too fast or too slow embryo cleavage rate has a negative impact on embryo development (Edwards et al., 1980; Kroener et al., 2015; Shebl et al., 2021). For example, it has been shown that fast growing embryos on Day 3 (>8 cells) have a higher rate of aneuploidy and an increased incidence of abnormal cleavage patterns and

are less likely to make blastocysts than 8-cell embryos (Kroener et al., 2015; Kong et al., 2016; Pons et al., 2019). However, once fast-growing embryos reach the blastocyst stage, their developmental potential is similar to 8-cell embryos (see also section 'Blastocyst stage (days 4–7)'). In contrast, concerning slow-developing embryos (<4 cells on Day 2, <8 cells on Day 3), there is clear evidence that these always perform worse and should only be used for transfer if better embryos are not available (Alikani et al., 2000; Thurin et al., 2005; Scott et al., 2007). These observations have been confirmed by embryo assessment using TLT (Meseguer et al., 2011; Montag et al., 2011; Herrero et al., 2013; Milewski et al., 2015).

Several studies using static observation have found speed of development to be predictive of live birth. In a prospective cohort study including 6252 Day-2 single embryo transfers, number of cells, the number of mononucleated cells per embryo and fragmentation rate were found to be significant predictors of live birth, with 4 cells and low (<10%) fragmentation having the highest live birth rate (Rhenman et al., 2015). In the most recent analysis of SART data including 28 878 fresh Day-3 embryo transfers, it was shown that for women at 34 years of age, the highest live birth rates were found after transfer of 8-cell embryos (24%), followed by >8 cell (23%), 7-cell (17%), 6-cell (8%), 5-cell (5%), and 4-cell (1%) embryos (Awadalla et al., 2022a). The 8-cell embryos with low degree of fragmentation (<10%) showed higher live birth rate compared to embryos with more than 10% fragmentation.

In addition, when looking at available evidence it should be considered that cell numbers on a specific day may be impacted by culture conditions and timing of assessments. It may also be challenging at times to distinguish between a cell and a large fragment. Obviously, assessment of Day-2 and Day-3 embryos by TLT permits more exact assessment timings, as well as detailed analysis of the developmental parameters over time, and the incidence of abnormal cleavages. For example, it is possible that some embryos with >8 cells on Day 3 are generated from trichotomous cleavages. This abnormal division can affect viability, but it is only detectable by TLT.

Fragmentation

A fragment can be defined as a membrane-bound extracellular cytoplasmic mass, often not including chromosomes. Fragments can vary in size and in distribution with different implications for the embryo (Alikani et al., 1999; Cecchele et al., 2022). The degree of fragmentation is difficult to evaluate, as it is first necessary to differentiate fragments from cells, and to consider the origin and to estimate the relative proportion of the embryo that is fragmented. One study found that a majority of blastomeres of <45 μm diameter in a Day-2 embryo and <40 μm diameter in a Day-3 embryo did not contain nuclei (Johansson et al., 2003). The impact of <10% fragmentation in Day-3 embryos on implantation rate has been found to be negligible (Alikani et al., 1999; Ebner et al., 2001; Van Royen et al., 2001; Holte et al., 2007; Racowsky et al., 2011), while, as discussed above, both earlier and several more recent and large studies, including TLT studies, have shown negative correlation with increasing fragmentation on live birth rates after early transfer (Rhenman et al., 2015; Ahlstrom et al., 2016; Awadalla et al., 2022b). Interestingly, a study by Ahlstrom et al., indicated that for Day-2 and Day-3 embryos, AI score correlated significantly with cell number and fragmentation score (Ahlström et al., 2023).

In addition, a correlation has been shown between the degree of fragmentation and the incidence of aneuploidy (Munné et al., 1995; Ziebe et al., 2003; Chavez et al., 2012).

Uneven cleavage and cell size

Uneven cellular cleavage, leading to unequal relative cell size, is commonly found in human embryos *in vitro* (Puissant et al., 1987). Unequal cell size has been defined as a 25% difference between the average diameter of the smallest cells compared to the average of the largest cells (Meseguer et al., 2011; Ziebe, 2013). Uneven cellular cleavage and its negative impact on pregnancy outcome for early transfer has been confirmed by several studies (Giorgetti et al., 1995; Ziebe et al., 1997; Hardarson et al., 2001; Racowsky et al., 2011), although some data are conflicting (Holte et al., 2007).

Interestingly, late-cleaving embryos have been reported to cleave more unevenly which, in turn, has been strongly correlated with an increased incidence of chromosomal errors (Hardarson et al., 2001; Shenoy et al., 2021), possibly due to uneven distribution of proteins, mRNA and mitochondria (Antczak and Van Blerkom, 1999).

It is important to consider that the relative cell sizes must be 'cell stage appropriate', i.e. assessed in relation to the number of cycles that cells have gone through. This means that the sister blastomeres representing the same cell cycle should be equally sized, i.e. only at the 2-, 4-, and 8-cell stage should all the cells be of the same size.

Multinucleation

Multinucleation has been correlated with a higher degree of fragmentation and decreased number of blastomeres on Days 2 and 3 (Van Royen et al., 2003), as well as with uneven cell size (Kligman et al., 1996; Hardarson et al., 2001; Sayed et al., 2022). The presence of multinucleation is generally considered abnormal, however the reported incidence varies greatly. The term 'multinucleation' can include different types of nucleation in one or more cells, including multiple (equally sized) nuclei, two nuclei (binucleation) and/or smaller size or micro nuclei (micronucleation). Most studies have not differentiated clearly between the different types, or in how many of the cells the condition is present, which may be a reason for some conflicting reports. For example, one study reported that 43% of patients had one or more embryo with multinucleation at the 2-cell stage, defined as ≥ 2 nuclei, which was reduced to 15% at the 4-cell stage (Balakier and Cadesky, 1997). Two other studies reported its occurrence in up to 87% of cycles, with 31–33% of the embryos affected at transfer (Jackson et al., 1998; Van Royen et al., 2003). Significantly slower development rates as well as lower implantation and live birth rates after early embryo transfer have been shown for embryos with multinucleation on Day 2 compared to mononucleated embryos (Ergin et al., 2014; Desch et al., 2017).

One recent TLT study, however, found that embryos that were binucleated at the 2-cell stage showed improved blastocyst formation rates and implantation rates, both compared to 'true' multinucleated embryos (≥ 3) and non-multinucleated embryos (Talbot et al., 2022). This shows the importance of distinguishing between the different types of nucleation during embryo assessment. Nucleation has shown to be a dynamic process, and the rate of multinucleation seen at the 2-cell stage is significantly reduced by the 4-cell stage (Aguilar et al., 2016; Balakier et al., 2016; Sayed et al., 2022; Talbot et al., 2022). It could also be that many of these embryos were binucleated but not 'true' multinucleated (≥ 3 nuclei) on Day 2, and should not be considered compromised, as discussed in the study by Talbot et al. (2022).

Evidence collected via TLT, where the cells can be scored in much more detail, has shown an incidence of 29–43% in multinucleation in early (2-cell stage) embryos with a significant impact

on implantation and live birth (Balakier et al., 2016; Goodman et al., 2016; Desch et al., 2017; Sayed et al., 2022). One study found an incidence of 6% multinucleated embryos with static scoring, compared to 23% using TLT (Ergin et al., 2014). Another study similarly found 7% and 35% using the two methods (Goodman et al., 2016).

In a further TLT study, it was shown that embryos with direct uneven cleavage or irregular chaotic divisions at the 2–5 cell stage showed a lower developmental potential. However, for those that did develop to the blastocyst stage, the presence of a single abnormality (multinucleation, reverse cleavage, irregular chaotic division, or direct uneven cleavage) at an early cell stage was not associated with aneuploidy when analysed at the blastocyst stage (Desai et al., 2018), while the presence of two or more abnormalities increased the risk of aneuploidy.

Other morphological features of Day-2 and Day-3 embryos

There is no conclusive evidence that embryos with apparent spatial disorganization, i.e. those that do not have the expected three-dimensional arrangement of blastomeres, should be considered abnormal (Ebner et al., 2012; Cauffman et al., 2014; Ebner et al., 2017; Desai and Gill, 2019).

Other morphological features, such as cytoplasmic granularity, membrane appearance and the presence of vacuoles can also be scored as part of the morphological assessment of Day-2 and Day-3 embryos (Magli et al., 2012). It is important to understand that these features can vary within and between cohorts.

Initiation of compaction

Compaction usually starts at the 8- to 16-cell stage. To be more precise, compaction spans the phase between the point in time when any two blastomeres of the multicellular embryo start to compact and the moment prior to the onset of blastocoel formation (Ciray et al., 2014). One study showed that almost 90% of embryos started compaction at the 8-cell stage or later (Iwata et al., 2014). Of these, 50% developed into good quality blastocysts, while for embryos that initiated compaction before the 8-cell stage, <20% became good quality blastocysts. Several other studies showed that beginning compaction on Day 3 can be a positive feature (Alikani et al., 2000; Skiadas et al., 2006; Le Cruguel et al., 2013; Aslan Öztürk et al., 2022). It is noteworthy that compaction on Day 2 is atypical and of unknown biological significance.

Consensus points

- Cleavage-stage embryo assessment should include cell number, grade and reason for the grade (e.g. 4-cell, grade 2, fragmentation), as previously agreed in the Istanbul Consensus (2011).
- Two-cell embryos on Day 1, 4-cell embryos on Day 2, and 8-cell embryos on Day 3, showing <10% fragmentation, mononucleation, and stage-specific cell size, should be prioritized in case of cleavage stage transfer or cryopreservation.
- There is no significant body of evidence to support an impact on implantation potential for cleavage stage embryos with atypical features such as spatial disorganization, vacuoles, cytoplasmic granularity, and zona abnormality, and these are therefore considered suitable for clinical use. However, extended culture of such embryos as a way of further selection for viability and evaluation should be considered.
- *Early cleavage*: The importance of assessing early cleavage for prediction of success rates has not been conclusively established. However, it may add information regarding other features such as binucleation/multinucleation and cell size.

Table 5. Ranking Scheme for Day-2 and Day-3 embryo transfer.

Feature	Top ranking	Intermediate ranking	Low ranking
Number of cells	4 cells on Day 2 or 8 cells on Day 3	>4 cells on Day 2 or >8 cells on Day 3	<4 cells on Day 2 or <8 cells on Day 3
Early cleavage	Early cleavage	No early cleavage	
Cell size	Cell stage specific	Not cell stage specific	
Fragmentation	None or minimal fragmentation (<10%)	10–25% fragmentation	>25% fragmentation
Multinucleation	No multinucleation at any cell stage	No multinucleation at 4 cell stage	Multinucleated at 4-cell stage
Abnormal cleavage	–	–	Direct cleavage DC2 (2- to 5-cell)
Compaction	Compaction from ≥8-cell stage	No compaction	Compaction before 8-cell stage
Recommendation	<ul style="list-style-type: none"> • De-prioritize Day 2/3 embryos with abnormal cleavage: direct cleavage DC1 (1- to 3-cell), irregular chaotic division or reverse cleavage, for transfer. • Extend culture of embryos with abnormal cleavage to blastocyst stage. 		

Assessment of early cleavage by TLT can be used to identify abnormal early cleavages such as direct cleavage, reverse cleavage and irregular chaotic division.

- **Fragmentation:** The relative degree of fragmentation was defined as: none or minimal (<10%), mild (<25%), or severe (>25%). The percent values are based on the cell equivalents, so for a 4-cell embryo, 25% fragmentation would equate to one blastomere in volume.
- **Numbers of blastomeres on Day 2/3:** The current expected observation for embryo development is 4 cells on Day 2 and 8 cells on Day 3. However, this can be influenced by the exact time of observation and culture conditions. It is recommended that the time of assessment is documented.
- **Cell size:** For embryos at the 2-, 4-, and 8-cell stages, blastomeres should be evenly sized. For all other cell stages, one would expect a cell stage appropriate size difference as the cleavage phase has not been completed.
- **Multinucleation:** True multinucleation (≥3 nuclei in one or several cells) is associated with decreased implantation potential and increased chromosome abnormality. Binucleation on Day 2, at the 4-cell stage, may not be necessarily a negative sign, but more evidence is needed. Laboratories should record the incidence and discriminate between binucleation, multinucleation and micronucleation in each embryo, and ideally, the nucleation status of each blastomere in each embryo. If available, multinucleation should be assessed using TLT.
- **Time-lapse technology:** Large datasets including timing of certain developmental events have been analysed to design algorithms to predict implantation and live birth. However, there is currently limited good quality evidence of better clinical outcomes following TLT embryo selection (Armstrong et al., 2019; Kieslinger et al., 2023). TLT allow assessment of kinetic variables such as rapid cleavage, direct cleavage, and reverse cleavage. These data have been used for deselection of embryos and it has been demonstrated that certain atypical cleavage patterns such as direct cleavage to three cells negatively affect embryo development. These events would in most cases be missed with static observations.
- **Compaction:** Based on a few studies, the start of compaction before 8 cells seems to negatively affect blastocyst formation, while compaction from 8 cells and onwards may be a positive indicator and could potentially be used as an additional selection tool at this stage.

Ranking cleavage-stage embryos

Different morphological features can reflect the overall quality of Day-2 and Day-3 embryos and the combination of those morphological features can be used to define a ranking order for transfer

or cryopreservation of Day-2 and Day-3 embryos. A proposed ranking scheme for Day-2 and Day-3 embryos is presented in Table 5.

5. Morula stage

When using TLT, the term morula refers to the 'end of the compaction process' (Ciray et al., 2014). Due to the variation in developmental speed and cellular complexity, there is a lack of well-defined temporal and morphological markers of morula development and viability for this stage (Coticchio et al., 2019). For an overview of all recommendations on morula stage assessment, see Table 6.

Timing of morula assessment and scoring

Accordingly, a morula would be the expected developmental stage if embryo scoring is done on Day 4 at 92 ± 2 hpi as recommended by the Istanbul Consensus (2011) (Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology, 2011).

The survey results showed that 24% of the respondents always apply the Istanbul Consensus (2011) recommendations related to the timing of assessment of Day-4 embryos (Supplementary Data SII, Fig. 3A).

However, TLT data have shown that there are considerable deviations in cleavage timings among a cohort of embryos of the same patient. At the extreme, a one-day delay or speed-up can be observed (Shebl et al., 2021), with neither scenario being necessarily associated with a worse treatment outcome.

Morphological features to consider for morula assessment

The survey results showed that 28% of the respondents always apply the Istanbul Consensus (2011) scoring criteria to score Day-4 embryos (Supplementary Data SII, Fig. 3B).

Timing of cavitation

Early cavitation of morulae is a good prognostic parameter related to better quality blastocysts with a higher potential to implant and higher ongoing pregnancy rates possibly due to a higher rate of euploidy (Hung et al., 2018). On the other hand, a delay in compaction and onset of cavitation was found to be associated with reduced blastocyst quality (Ivec et al., 2011; Desai et al., 2014) and reduced likelihood of live birth (Fisheh et al., 2018).

Number of cells

Quality assessment at 92 ± 2 hpi usually takes both cell number and degree of compaction into consideration (Alikani et al., 2000; Tao et al., 2002; Feil et al., 2008; Ebner et al., 2009; Fabozzi et al., 2016). It has been found that the more cells and in particular the

Table 6. Overview of all evidence and recommendations for morula stage/Day-4 embryo assessment.

Feature	Summary of review findings				Considerations	Recommendation
	Atypical patterns	Embryo quality and development potential	Ploidy	Implantation rate		
Timing of cavitation	Early cavitation	N/R	Association with higher euploidy rate Very low ⊕○○○ 1 observational study (Hung et al., 2018)	Association with higher implantation rate Very low ⊕○○○ 2 observational studies (Hung et al., 2018; Rienzi et al., 2019)	Association with higher ongoing pregnancy rate Very low ⊕○○○ 2 observational studies (Hung et al., 2018; Rienzi et al., 2019)	Day-4 embryos showing full compaction or early cavitation should be prioritized in case of Day-4 transfer or cryopreservation.
	Delay in compaction	Association with lower blastocyst quality Very low ⊕○○○ 2 observational studies (Ivec et al., 2011; Desai et al., 2014)	Contradictory results: No clear association with aneuploidy rate Very low ⊕○○○ 1 observational study (Minasi et al., 2016) Association with higher euploidy rate Very low ⊕○○○ 1 observational study (Campbell et al., 2013)	No clear association with implantation rate Very low ⊕○○○ 1 observational study (Montjean et al., 2021)	No clear association with live birth rate Very low ⊕○○○ 1 observational study (Montjean et al., 2021)	
Number of cells	More compacting cells on Day 4	Correlation with blastocyst formation rate Very low ⊕○○○ 2 observational studies (Ebner et al., 2009; Iwata et al., 2014)	N/R	N/R	N/R	Accurate evaluation of cell number is impossible once the majority of blastomeres is involved in the compacting mass, and the focus is placed on the proportion of cells involved in compaction.
Degree of compaction	Partly compacted embryos (excessive fragmentation, large number of excluded cells, self-cavitation of blastomeres)	Association with lower blastocyst formation rate and blastocyst quality Low ⊕○○○ 5 observational studies (Alikani et al., 2000; Ebner et al., 2009; Lagalla et al., 2017; Coticchio	N/R	N/R	Association with lower live birth rate Very low ⊕○○○ 1 observational studies (Coticchio et al., 2021)	Embryos with partial compaction can form blastocysts and should be considered for clinical use. Extended culture of these embryos for further evaluation should be considered.

(continued)

Table 6. Continued

Feature	Summary of review findings				Considerations		Recommendation
	Atypical patterns	Embryo quality and development potential	Ploidy	Implantation rate	Live birth rate		
Vacuolization	<p>Vacuole formation around compaction</p> <p>Compaction of vacuolized blastomeres</p>	<p><i>et al., 2021; Parriego et al., 2024</i>)</p> <p>Association with lower blastocyst formation rate and blastocyst quality</p> <p>Very low ⊕○○○ 2 observational studies (<i>Mayer et al., 2018; Chen et al., 2019</i>)</p>	N/R	N/R	<p>Association with lower ongoing pregnancy rate and live birth rate</p> <p>Very low ⊕○○○ 2 observational studies (<i>Feil et al., 2008; Mayer et al., 2018</i>)</p>	<p>No correlation has been found between the occurrence of vacuoles and patient parameters like age or baseline hormonal profile</p>	<p>Spontaneous vacuole formation around compaction was found to be a negative predictor for embryo development.</p>
Cleavage dynamics	<p>Blastomere exclusion/extrusion</p>	<p>N/R</p> <p>Contradictory results: Higher aneuploidy in excluded cells</p> <p>Very low ⊕○○○ 1 observational study (<i>Lagalla et al., 2017</i>)</p> <p>Ploidy correlation with excluded cells</p> <p>Very low ⊕○○○ 1 observational study (<i>Parriego et al., 2024</i>)</p>	N/R	N/R	<p>Association with lower live birth rate</p> <p>Very low ⊕○○○ 2 observational studies (<i>Coticchio et al., 2021; Hür et al., 2023</i>)</p>	<p>Blastomere exclusion/extrusion at morulae stage is likely to be associated with abnormalities in the eliminated cells.</p>	<p>Normally cleaving embryos result in euploid blastocysts less frequently than their irregular cleaving counterparts.</p>

N/R, not reported.

Table colour code: Green: the embryo can be clinically used; Yellow: the embryo could be used with cautionary considerations; Red: the embryo is not considered suitable for clinical use.

more compacting cells a Day-4 embryo shows the better its chance of forming a blastocyst on Day 5 (Ebner et al., 2009; Iwata et al., 2014).

Since accurate evaluation of cell number is impossible once the majority of blastomeres is involved in the compacting mass, focus is placed on the proportion of cells involved in compaction. In principle, partly (PCM) and fully (FCM) compacted morulae can be distinguished. The former group is characterized by a certain loss of embryonic mass either due to extensive cytoplasmic fragmentation or blastomere elimination. If the observed loss is substantial, further development to blastocyst (Alikani et al., 2000; Ebner et al., 2009; Lagalla et al., 2017; Coticchio et al., 2021) and formation of good quality blastocysts (Ebner et al., 2009; Coticchio et al., 2021) will be affected, both of which could be associated with a lower live birth rate (Coticchio et al., 2021).

Other morphological features

Beyond the degree of compaction, some studies have also considered detrimental morphological features such as: excessive fragmentation, multiple excluded cells, 'self-cavitation' of blastomeres and vacuolization for morphological assessment of Day-4 embryos (Alikani et al., 2000; Feil et al., 2008; Ivec et al., 2011; Fabozzi et al., 2016). Of note, the first three abnormalities would reflect PCM, which implies that vacuolization is the only abnormality that could be taken into consideration for quality assessment purposes. Indeed, spontaneous vacuole formation around the time of compaction was found to be a negative predictor of blastulation and top-quality blastocyst formation rates (Mayer et al., 2018; Chen et al., 2019), ongoing pregnancy rates (Feil et al., 2008; Mayer et al., 2018) and live birth rates (Mayer et al., 2018).

Recent TLT studies further shed some light on the phenomenon of blastomere loss around the morula stage (Lagalla et al., 2020; Coticchio et al., 2021). Two types of cleavage dynamics were identified, both of which were responsible for the elimination of blastomeres but differed in timing. One was the exclusion of blastomeres from the outset and the other was characterized by the extrusion of cells after full compaction had already occurred. The occurrence of the two phenomena together had the worst prognosis for live birth (Coticchio et al., 2021; Hur et al., 2023).

Blastomere exclusion/extrusion at morula stage is likely to be associated with abnormalities in the eliminated cells. It has been shown that excluded cells show E-cadherin (a key cell adhesion protein) expression profiles that are different from the expected membrane-localized pattern (Alikani, 2005). The degree to which failed compaction or blastomere loss (Zhu et al., 2021) at compaction reflects perturbations in key events in compaction and cell polarization of the morula (e.g. apical F-actin and PAR complex accumulation) remained speculative until recently, when it became evident that contractile forces of cells play a key role in the compaction process. The fact that embryos that fail to compact or exclude cells exhibit lower surface tension suggests that weak cell contractility is the causative phenomenon (Firmin et al., 2024). In relation to partial compaction, other studies reported 'abnormal divisional behaviour' such as karyokinesis without cytokinesis or signs of degeneration (Zhan et al., 2016). The appearance of apoptotic nuclei following compaction further suggests that programmed cell death may play a role in eliminating affected blastomeres (Chatzimeletiou et al., 2005).

A more detailed annotation of the TLT sequences revealed that in comparison to FCM all patterns of PCM not only show a higher rate of irregular and asymmetric cleavage (Coticchio et al., 2021) but also an evident delay in development starting with

pronuclear fading (Lagalla et al., 2020; Coticchio et al., 2021; Hur et al., 2023). In particular, highly dynamic biological processes such as compaction and blastulation were deferred (Lagalla et al., 2020; Coticchio et al., 2021; Ezoe et al., 2023).

A hierarchical classification model has found morula formation (tM) within an optimal range (81.3–96.0 hpi) to be one of the strongest predictors of blastocyst formation (Motato et al., 2016). Similarly, a multivariate analysis has shown that tM was the only morphokinetic parameter that correlated with live birth rate after euploid blastocyst transfer (Rienzi et al., 2019).

While some studies showed no correlation between tM or starting blastulation (tSB) and aneuploidy (Minasi et al., 2016) others found a delayed initiation of compaction (tSC) in complex aneuploid embryos (Campbell et al., 2013).

There is evidence that PCM following irregular cleavages can develop into euploid blastocysts (Zhan et al., 2016; Lagalla et al., 2017). Those cells excluded from the morulae were shown to have a high rate of aneuploidy and degraded DNA (Lagalla et al., 2017). This, together with reduced aneuploidy rate in biopsied TE cells of the associated blastocyst, suggests that a self-check mechanism may reduce the relative abundance of aneuploid cells.

On the other hand, a recent study showed a high ploidy correlation between excluded cells and TE cells, suggesting that cell exclusion might be a consequence of compromised embryo development regardless the chromosomal constitution of excluded cells (Parriego et al., 2024).

Consensus points

- Day-4 embryos showing full compaction or early cavitation should be prioritized in case of Day-4 transfer or vitrification.
- Embryos with partial compaction can form blastocysts and should be considered for clinical use. Extended culture of these embryos for further evaluation should be considered.

A proposed ranking scheme for morulae is presented in Table 7.

6. Blastocyst stage (Days 4–7)

Embryo culture to the blastocyst stage is routine in clinical embryology encompassing Days 4 to 7 and represents a significant shift in practice since the Istanbul Consensus was first published in 2011.

The survey results indicate that only 27% of the respondents follow the Istanbul Consensus (2011) recommendations on the timing and criteria for scoring blastocysts. The Gardner grading system (Gardner and Schoolcraft, 1999), remains the most common scheme utilized clinically, according to the survey results (63% of respondents) (Supplementary Data SII, Fig. 1D). Re-evaluation and modification of the Gardner grading system was to be expected and this has indeed occurred (Veck and Zaninovic, 2003; Cuevas Saiz et al., 2018; Hammond et al., 2020; Pierson et al., 2023), and 30% of respondents indicated using an additional grade (either 'D' or 'X') or the term 'non-classifiable' to denote blastocysts considered unsuitable for clinical use.

AI has been applied to both consecutive images of embryo development obtained through time-lapse (Khosravi et al., 2019; Tran et al., 2019; Berntsen et al., 2022; Illingworth et al., 2024), and to static images of blastocysts (Bormann et al., 2020; Chavez-Badiola et al., 2020; Diakiw et al., 2022), in an attempt to improve the ability to identify the most viable embryo in a cohort, while reducing the intra- and inter-operator variation associated with subjective evaluation of blastocysts using the grading systems

Table 7. Ranking for selection of morulae with similar hours post-insemination (hpi).

Feature	Top ranking	Intermediate ranking	Low ranking
(Early) cavitation	Yes	No	No
Compaction	FCM	PCM	No compaction Compacting embryo with ≥8 cells PCM with significant cytoplasmic loss
Morphology	No vacuoles	No to minor vacuolisation	Heavy vacuolisation
Recommendation:	Extend culture to blastocyst for embryos with atypical morphological features: self-cavitation of blastomeres, <50% compacted embryo, ≤8 cells without compaction, excessive fragmentation, widespread vacuoles.		

FCM, Fully compacted morulae; PCM, partially compacted morulae.

discussed. Interestingly, a recent paper by Ezoë *et al.*, indicated that AI score was tightly coupled to the morphological aspects of the Gardner grading system (Ezoë *et al.*, 2022b). AI holds great promise to augment embryologist assessment of the blastocyst (Fitz *et al.*, 2021; Sawada *et al.*, 2021), but should not yet be considered as a replacement for conventional assessment (Illingworth *et al.*, 2024). The survey results showed that only 14% of the respondents make use of AI mainly for embryo assessment in TL videos (in 71% of cases) (Supplementary Data SII, Fig. 6C).

For an overview of all recommendations on blastocyst assessment, see Table 8.

Timing of blastocyst scoring

The recommended timing by the Istanbul Consensus (2011) for static observation of Day-5 embryos is 116h±2 hpi (Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology, 2011). However, formation and expansion of a blastocoel cavity in embryos leading to a live birth occurs over a wide timeframe, from as early as Day 4 (98.4±0.4 hpi) to the 'typical' timing of Day 5 (112.4±0.1 hpi) or delayed until Day 6 (131.6±0.1 hpi) or Day 7 (151.2±0.5 hpi) (Coticchio *et al.*, 2023). Maintaining a standardized window for embryo assessment can be beneficial for benchmarking, establishing and monitoring KPIs, although this should be balanced against workflow needs, particularly when TLT is not available (Figure 1). In terms of timing of assessment, even if daily assessment timings cannot be consistent, blastocysts within a cohort can be compared for developmental stage as well as morphology to aid selection, while being mindful of reports that faster developing embryos, at each stage of development, have greater potential for implantation and birth, than their slower counterparts (Campbell *et al.*, 2022b).

Morphological features to consider for blastocyst assessment

Day of blastocyst formation

Developmental speed is directly correlated with blastocyst viability: slower growing blastocysts have lower implantation rates (Shebl *et al.*, 2021). While blastocysts developing according to the expected timeline have high implantation rates when transferred during a fresh cycle (Shebl *et al.*, 2021), slow growing blastocysts

may miss the window of implantation, a problem that is partially alleviated with blastocyst vitrification and transfer in a frozen cycle (Day 5 vs Day 6, RR 1.74 (95% CI 1.37–2.20) for fresh transfer and 1.38 (95% CI 1.23–1.56) for frozen embryo transfer (FET)) (Bourdon *et al.*, 2019), particularly for Day-6 blastocysts that were at the morula stage on Day 5 (Tannus *et al.*, 2019). Day-4 blastocysts, although rare, display a very high implantation rate in FET cycles (Coticchio *et al.*, 2023).

Live birth rates for untested blastocysts frozen on Day 6 are lower than those frozen on Day 5 (Bourdon *et al.*, 2019; Yerushalmi *et al.*, 2021; Coticchio *et al.*, 2023); and this difference persists with the transfer of euploid blastocysts (Tiegs *et al.*, 2019; Zhan *et al.*, 2020; Cimadomo *et al.*, 2022b; Lane *et al.*, 2022). Day-7 blastocysts, which may represent 5–10% of all useable blastocysts (Hammond *et al.*, 2018), have higher rates of aneuploidy and lower implantation rates compared to Day-5 and Day-6 euploid blastocysts (Tiegs *et al.*, 2019; Cimadomo *et al.*, 2022b; Lane *et al.*, 2022). Nonetheless, healthy live births can be obtained with Day-7 blastocysts and these embryos may be of particular importance for patients with few embryos available (Du *et al.*, 2018) or with advanced maternal age (Abdala *et al.*, 2023). Survey results indicated that a small minority (16%) of the respondents perform some fresh Day-7 blastocyst transfers, while most others (49%) transfer Day-7 blastocysts in frozen embryo transfer cycles.

Degree of expansion and ICM/TE grade

Implantation potential according to the Istanbul Consensus (2011) scoring system is related to expansion stage and ICM/TE grade, though the relative importance of each remains to be fully resolved. The difference between ICM/TE grades A and B appears marginal, whereas grade C is considered non-useable by 44% of respondents. The remaining respondents use a modified Gardner grade or the term 'non-classifiable' and consider blastocysts with grade C ICM or TE as useable. This marked difference in clinical practice indicates lack of consensus, an observation further supported by the finding that 8 of 10 respondents indicated that a universally accepted term for non-useable blastocysts is needed.

Fresh untested blastocyst transfers represent a significant proportion of treatment cycles and have helped establish the relative importance of blastocyst characteristics. Multivariate




Figure 1. Time windows chart: this figure provides an example of suggested timings for assessment, to maximize the chance of observing the developing embryo at specific stages. In this example, IVF/ICSI is performed at 3:00 pm. hpi: hours post insemination.

analysis accounting for expansion stage, ICM grade and TE grade shows that grade of TE is the strongest predictor of live birth (Ahlström et al., 2011; Hill et al., 2013; Thompson et al., 2013; Ebner et al., 2016; Bakkensen et al., 2019; Pons et al., 2023), followed by degree of expansion (Thompson et al., 2013; Du et al., 2016; Subira et al., 2016; Bakkensen et al., 2019). Few blastocysts with grade 'C' ICM or TE were included in these studies; notably one study found Grade 'C' ICM was associated with lower live birth rate (Subira et al., 2016). In general, expanded blastocysts with higher grade TE are associated with higher live birth rates in fresh transfers (Zou et al., 2023). Similarly, in a multivariate analysis of over 2000 fresh blastocyst transfers, one study showed that both expansion stage and TE grade were associated with the probability of live birth (Storr et al., 2019). The impact of ICM grade on outcome is less clear. While ICM grade may be associated with pregnancy loss (Van den Abbeel et al., 2013), and birthweight (Licciardi et al., 2015), further evidence is needed to establish definitive links. Blastocysts showing marked signs of

degeneration or without clearly discernible ICM may sporadically produce live births, but pertinent evidence is anecdotal (Kovacic et al., 2004).

Predictive features of untested fresh and frozen blastocysts compare favourably. TE grade was the most common variable associated with live birth from frozen blastocysts (Honma et al., 2012; Ahlström et al., 2013; Chen et al., 2014), followed by expansion stage (Ahlström et al., 2013). None of these studies found an association between ICM grade and implantation, though similar to studies with fresh blastocysts, grade 'C' ICM was not well represented in frozen embryo transfer cycles. Of note and in contrast to fresh transfers where only Day-5 embryos were transferred, none of the studies controlled for day of blastocyst formation in the multivariate analysis, thus limiting their applicability for using stage/grade when ranking slower growing blastocysts.

Though most studies have found that TE grade has the highest correlation with live birth, at least one multivariate analysis

Table 8. Overview of all evidence and recommendations on blastocyst assessment.

Feature	Summary of review findings				Recommendation	
	Atypical patterns	Embryo quality and development potential	Ploidy	Implantation rate		
Day of blastocyst formation	Slow blastocyst (Day 5 vs Day 6)	N/R	N/R	Association with lower implantation rates Very Low ⊕○○○ 1 observational study (Shebl <i>et al.</i> , 2021)	Association with lower live birth rate Low ⊕⊕○○ 3 observational studies (Bourdon <i>et al.</i> , 2019; Yershalmi <i>et al.</i> , 2021; Coticchio <i>et al.</i> , 2023)	Slow growing blastocysts may miss the window of implantation, a problem that is partially alleviated with blastocyst vitrification and transfer in a frozen cycle.
	Day 7 blastocysts	N/R	Association with higher aneuploidy rates Low ⊕⊕○○ 3 observational studies (Tiegs <i>et al.</i> , 2019; Cimadomo <i>et al.</i> , 2022b; Lane <i>et al.</i> , 2022)	Association with lower implantation rates Low ⊕⊕○○ 3 observational studies (Tiegs <i>et al.</i> , 2019; Cimadomo <i>et al.</i> , 2022b; Lane <i>et al.</i> , 2022)	Association with lower live birth rate Low ⊕⊕○○ 1 review and 3 observational studies (Hammond <i>et al.</i> , 2018; Tiegs <i>et al.</i> , 2019; Cimadomo <i>et al.</i> , 2022b; Lane <i>et al.</i> , 2022)	Speed of development is one of the primary indicators of blastocyst potential and should be used for ranking of blastocysts for transfer Day-7 blastocysts can be viable and could be considered for clinical use.
Grade	Degree of expansion	N/R	Contradictory results: Association with higher aneuploidy rates Very low ⊕○○○ 3 observational studies (Campbell <i>et al.</i> , 2013; Huang <i>et al.</i> , 2019; Cimadomo <i>et al.</i> , 2022b) No clear association with aneuploidy rate Very low ⊕○○○ 3 observational studies (Kramer <i>et al.</i> , 2014; Yang <i>et al.</i> , 2014; Rienzi <i>et al.</i> , 2015)	N/R	Association with higher live birth rate Very low ⊕○○○ 6 observational studies (Ahlström <i>et al.</i> , 2013; Thompson <i>et al.</i> , 2013; Du <i>et al.</i> , 2016; Subira <i>et al.</i> , 2019; Bakkensen <i>et al.</i> , 2019; Storr <i>et al.</i> , 2019)	Degree of expansion is one of the primary indicators of blastocyst potential and should be used for ranking of blastocysts for transfer.
	ICM/TE grade	N/R	Association with aneuploidy rate Moderate ⊕⊕⊕○ 1 systematic review and 5 observational studies (Huang <i>et al.</i> , 2021; Zou <i>et al.</i> , 2022; Bamford <i>et al.</i> , 2023; Barnes <i>et al.</i> , 2023; Hori <i>et al.</i> , 2023; Kato <i>et al.</i> , 2023)	Contradictory results: ICM grade associated with implantation Very low ⊕○○○ 5 observational studies (Irani <i>et al.</i> , 2017; Zhao <i>et al.</i> , 2018; Nazem <i>et al.</i> , 2019; Abdala <i>et al.</i> , 2022; Zhang <i>et al.</i> , 2022) No clear association of ICM grade with implantation rate Very low ⊕○○○	TE grade is the strongest predictor of live birth rates Low ⊕⊕○○ 10 observational studies (Ahlström <i>et al.</i> , 2011; Honma <i>et al.</i> , 2012; Ahlström <i>et al.</i> , 2013; Hill <i>et al.</i> , 2013; Thompson <i>et al.</i> , 2013; Chen <i>et al.</i> , 2014; Ebner <i>et al.</i> , 2016; Bakkensen <i>et al.</i> , 2019; Storr <i>et al.</i> , 2019; Pons <i>et al.</i> , 2023)	Trophectoderm is one of the primary indicators of blastocyst potential and should be used for ranking of blastocysts for transfer. Grade C blastocysts can be viable and could be considered for clinical use.

(continued)

Table 8. Continued

Feature	Summary of review findings				Considerations	Recommendation
	Atypical patterns	Embryo quality and development potential	Ploidy	Implantation rate		
Chromosomal status	Aneuploid	Association with lower embryo quality very low ⊕○○○ 3 observational study (Capalbo et al., 2014; Minasi et al., 2016; Kato et al., 2023)	N/R	3 observational studies (Honma et al., 2012; Ahlström et al., 2013; Chen et al., 2014) N/R	N/R	Identifying embryos at highest risk of being chromosomally abnormal is not a diagnostic approach but rather could be perceived as a mean to identify those blastocysts with greatest probability of being aneuploid and hence candidates for biopsy and genetic analysis.
Cytoplasmic strings	Presence of cytoplasmic strings	Association with higher blastocyst quality Very low ⊕○○○ 1 observational study (Ma et al., 2022)	N/R	Association with higher implantation rate Low ⊕⊕○○ (Ebner et al., 2020; Ma et al., 2022; Eastick et al., 2021; Eastick et al., 2023a; Joo et al., 2023)	N/R	The utility of cytoplasmic strings presence as an independent indicator for ranking is unknown.
Spontaneous collapse	Spontaneous collapse	Association with lower blastocyst quality Very low ⊕○○○ 1 observational study (Cimadomo et al., 2022a)	Association with euploidy rate Low ⊕⊕○○ 1 meta-analysis of 3 observational studies (Bickendorf et al., 2023)	No clear association with implantation potential Very low ⊕○○○ 2 observational studies (Sciorio et al., 2020; Cimadomo et al., 2022a)	Contradictory results: Association with lower ongoing pregnancy rate Low ⊕⊕○○ 1 meta-analysis of 5 observational studies (Bickendorf et al., 2023) No clear association with live birth rate Low ⊕⊕○○ 1 meta-analysis of two observational studies (Bickendorf et al., 2023)	The significance of spontaneous collapse on pregnancy outcomes is unclear.
ICM	Presence of 2 ICM Potential complication Very low ⊕○○○ 2 observational studies (Payne et al., 2007; Noli et al., 2015)	N/R	N/R	N/R	N/R	Blastocysts with 2 ICM indicating potential monozygotic twinning should not be transferred without thorough patient counselling. Non-viable blastocysts should be graded as 'D' as opposed to 'C' based on degenerative features or absence of a distinct ICM.
	Absence of ICM	N/R	N/R	Association with lower implantation rate Very low ⊕○○○ 1 observational study (Kovacic et al., 2004)	Association with lower live birth rate Very low ⊕○○○ 1 observational study (Kovacic et al., 2004)	Transfer of blastocysts without ICM may lead to abnormal pregnancy or pregnancy loss.

ICM, inner cell mass; N/R, not reported; TE, trophoctoderm.
Table colour code: Green: the embryo can be clinically used; Yellow: the embryo could be used with cautionary considerations. Red: the embryo is not considered suitable for clinical use.

found that the grade of the ICM is the variable most commonly associated with implantation (Irani et al., 2017). However, most of the studies only found an association with grade 'C' ICM, not between grade 'A' and 'B' (Zhao et al., 2018; Nazem et al., 2019; Abdala et al., 2022; Zhang et al., 2022). Some of these studies also found an association with TE grade (Zhao et al., 2018; Nazem et al., 2019) and expansion stage (Abdala et al., 2022). A recent study developed a composite blastocyst score where day, expansion stage, TE and ICM grades were all significantly associated with a clinical pregnancy, and blastocyst day had the largest impact, followed by ICM grade, expansion and TE grade (Zhan et al., 2020). It is acknowledged that while assessing the grade of the TE is relatively straight forward, assessing the ICM can be more problematic depending on its position and shape, and hence reflects the difficulties in differentiating between A and B grades of ICM in some blastocysts.

Early in the clinical application of blastocyst culture, a threshold for blastocyst useability was set at Gardner 3BB when slow freezing and variable cryosurvival influenced the decision (Langley et al., 2001). Since the adoption of vitrification and PGT-A, several studies indicate that presumably low-grade blastocysts classified as low viability (e.g. grade C) can produce healthy live births, albeit at greatly reduced rates (Morbeck, 2017; Kemper et al., 2021). Similar to Day-7 blastocysts, these low-grade blastocysts may be useful for patients with few available embryos (Cimadomo et al., 2022b). These changes to the definitions of usable blastocysts have raised important clinical and ethical questions: has the lower limit of viability been established and, if so, can these embryos be discarded where laws forbid destruction of human embryos? A framework for defining 'developmentally incompetent' preimplantation embryos has been developed to address this unique and important area of clinical practice (Cimadomo et al., 2021).

Abnormal chromosomal status

Human embryos with abnormal chromosomal status can develop as evidenced by the fact that specific trisomies are compatible with the formation of high scoring blastocysts, and some, such as trisomy 21, can go to term (Forman et al., 2013; Savio Figueira Rde et al., 2015). Importantly, blastocysts with abnormal chromosomal status will exhibit cellular stress, through which their transcriptome, proteome and metabolome will be affected, thereby compromising their physiology and development.

A relationship between blastocyst morphology and aneuploidy following TE biopsy was initially inferred by a retrospective observational study (Capalbo et al., 2014), which determined an incidence of aneuploidy of 6.8%, 15.2%, 17.4%, and 27.5% in excellent, good, average, and poor quality embryos, respectively, in women >35 years old. Significantly, in blastocysts where both ICM and TE were abnormal, there was a doubling in the frequency of aneuploidy. Another case series study with analysis of 1730 embryos reported that euploid blastocysts were characterized with high scoring ICM and TE, as well as a high degree of expansion, and a shorter time to the initiation of blastocoel formation (Minasi et al., 2016). Similarly, an analysis of 3573 blastocysts showed that euploidy was correlated with the Gardner grade but did not report the relative contributions of the grading to ploidy (Kato et al., 2023).

Using time-lapse to determine the timing of blastocyst formation (reflected in the expansion stage), it was observed that kinetics and rate of embryo expansion are related to aneuploidy risks (Campbell et al., 2013; Huang et al., 2019; Cimadomo et al., 2022b). However, other groups failed to confirm these findings (Kramer

et al., 2014; Yang et al., 2014; Rienzi et al., 2015). More recently, AI has been applied to analysing embryo morphology correlation with blastocyst euploidy rates (Huang et al., 2021; Zou et al., 2022; Bamford et al., 2023; Barnes et al., 2023; Hori et al., 2023; Kato et al., 2023) with promising results. Interestingly, AI score appears closely associated with the Day-5 Gardner grade in euploid blastocysts (Kato et al., 2023). While certain aspects of blastocyst morphology and specific AI have been able to identify those embryos at highest risk of being chromosomally abnormal, the approach lacks diagnostic accuracy. However, these methods could be used to identify those blastocysts with greatest probability of being aneuploid and hence candidates for biopsy and genetic analysis.

Spontaneous collapse

A benefit of time-lapse culture is the ability to assess poorly studied blastocyst features such as spontaneous blastocoel collapse. Approximately one in four blastocysts show spontaneous collapse and re-expansion and even fewer have more than one collapse (Marcos et al., 2015). The significance of a spontaneous collapse for ongoing pregnancy or live birth is unclear (Marcos et al., 2015; Bodri et al., 2016; Sciorio et al., 2020), though most evidence suggests a negative impact. Blastocysts that collapse are more likely to be aneuploid; however, some reports indicate a history of collapse does not affect euploid embryo implantation (Cimadomo et al., 2022a; Bickendorf et al., 2023).

Cytoplasmic strings

Cytoplasmic strings are dynamic structures connecting TE and ICM cells and are involved in cellular communication (Salas-Vidal and Lomelí, 2004). Appearing in 55–85% of expanded, transferred blastocysts, cytoplasmic strings are positively associated with implantation (Ebner et al., 2020; Eastick et al., 2021; Ma et al., 2022; Eastick et al., 2023a; Joo et al., 2023). Similar to blastocyst grading in general, assessment of cytoplasmic strings has fair to moderate inter- and intra-observer agreement (Eastic et al., 2023b). Though strings are associated with higher blastocyst quality (Ma et al., 2022), the utility of their inclusion as an independent predictor of viability for ranking is unclear. While the presence of strings is significantly associated with clinical pregnancy when controlling for degree of expansion and ICM/TE grade (Eastic et al., 2023b), the multivariate analysis did not account for day of blastocyst formation.

Other morphological features

The presence of two ICMs in one blastocyst is a rare occurrence and warrants careful consideration. Monozygotic twinning is a complication more common following assisted reproductive technologies with significant risks to the offspring and the mother (Vitthala et al., 2009; Hviid et al., 2018; Busnelli et al., 2019; Kadam et al., 2023). Since few case reports exist of blastocysts with two ICMs in vitro (Veck and Zaninovic, 2003; Payne et al., 2007; Noli et al., 2015), splitting of the ICM is unlikely to occur until after embryo transfer. Given the risks to the offspring and the mother, clinics may consider having a policy to not use blastocysts with suspected two or more ICM. Alternatively, when two ICM are visible prior to transfer, clinics should have a policy whereby the medical team is notified to allow for proper patient counselling.

Several other features beyond traditional morphology may also be used in ranking blastocysts. While many reports correlate early embryo developmental features with blastocyst implantation, most do not account for blastocyst morphology in the statistical analysis. The only pre-compaction variable associated with

blastocyst live birth, when accounting for blastocyst quality, is the number of cells on Day 3, where slow cleaving embryos (<7 cells) have reduced implantation rates when transferred at the blastocyst stage (Wu et al., 2020; Zhao et al., 2020). Utility of this finding is uncertain, however, since it would only be applied when selecting between two blastocysts with similar Day/ stage/grade.

Consensus points

- Ultimately, the goal of blastocyst grading is ranking for order of use.
- The Gardner grading system for blastocyst scoring should be used (Table 9; Supplementary Data SIV, Fig. 1). This system is distinguished from the prior Consensus grading by using letters for the ICM/TE grades and adding additional expansion stages (e.g. hatched blastocyst).
- Non-viable blastocysts should be graded as 'D' as opposed to 'C' based on degenerative features or absence of a distinct ICM.
- The common features that are clearly associated with implantation potential include day of blastocyst formation (Days 4–7), stage of expansion (3, 4, 5, 6), and grade of ICM (A, B, C) and TE (A, B, C).
- Blastocysts with grade C ICM and/or TE and Day 7 blastocysts can be viable and could be considered suitable for clinical use.
- Blastocysts with two ICM indicating potential for monozygotic twinning should not be transferred without thorough patient counselling.
- Assigning relative importance of each variable requires systematic multivariate analysis with a large dataset and is further complicated when assessing fresh versus frozen untested and euploid blastocysts.

7. Duration of embryo culture and frequency of assessments: safety versus effectiveness

The Istanbul Consensus (2011) offers a broad spectrum of morphological parameters for oocyte and embryo assessment. In laboratories using TLT-equipped incubators, continuous culture allows flexibility in the frequency and level of detail of embryo evaluation, without disturbing the culture conditions. In laboratories performing static observations, however, the frequency of embryo assessment should be determined considering factors such as the type of incubators used (bench top or big box), the type of culture medium (single or sequential), the use of isolettes for embryo handling, and the duration of embryo culture (cleavage or blastocyst stage). The aim is to strike an optimal balance between acquiring the desired information on developing embryos and minimizing the disturbance of the culture conditions (Swain, 2014; Wale and Gardner, 2016; ESHRE Working group on Time-lapse technology et al., 2020).

Some ART centres still combine cleavage and blastocyst stage embryo transfers, as shown in our survey (Supplementary Data SII, Fig. 4). The duration of embryo culture, embryo morphology assessment and embryo transfer policy, whether for fresh or frozen embryos, should primarily aim for the fastest, safest and most economically sustainable way to achieve the goal of fertility treatment. The choice of assessment methods, level of detail, and the duration and frequency of monitoring of embryo development under *in vitro* conditions should therefore be tailored to the available laboratory equipment.

Current practice of cleavage stage versus blastocyst transfer

Our survey showed that the blastocyst stage is commonly used in ART centres for performing embryo transfer. Fewer than 2% of ART centres did not perform blastocyst transfer at all while 17.4% performed blastocyst transfer nearly exclusively (in >95% of cycles) (Supplementary Data SII, Fig. 4A).

Interestingly, Day-2 and Day-3 embryo transfer were not practiced at all in 44% and 8% of ART centres, respectively. On the other hand, only 2–3% of ART centres exclusively practiced cleavage stage embryo transfer with 2.2% performing transfers on Day-3 and 0.7% on Day-2 (Supplementary Data SII, Fig. 4A).

Moreover, cryopreservation of blastocysts predominates over cleavage stage embryos. More than 50% of the respondents reported that embryos are exclusively cryopreserved at the blastocyst stage, while in the remaining cases mostly a combination of cryopreservation of Day-3 and Day-5/6 embryos is performed (Supplementary Data SII, Fig. 4B). Day-2 and Day-4 embryos are never cryopreserved by roughly 75% of ART centres (Supplementary Data SII, Fig. 4B). A similar trend with a higher percentage of blastocyst (73.9%) over cleavage stage (26.1%) frozen transfers can be found in the ESHRE report for 2018 (Wyns et al., 2022).

The transfer of Day-4 embryos occurred in <25% of the transfer cycles according to 36.3% of the respondents and only 19.9% of the respondents reported that they cryopreserve Day-4 embryos in <25% of the transfer cycles (Supplementary Data SII, Fig. 4). It is not clear whether the reason for the use of Day 4 embryos is the earlier development of the blastocyst or the earlier scheduling of the day of transfer or cryopreservation at the convenience of the patient or the centre.

Reasons for increasing use of extended embryo culture

Several factors have contributed to the increasing use of blastocyst transfer. There is consistent evidence from a multitude of studies showing higher pregnancy and live birth per transfer using fresh blastocyst transfer, with this observation being more prominent in good prognosis patients (Practice Committee of the American Society for Reproductive Medicine, 2018). However, a retrospective analysis of more than 100 000 IVF/ICSI cycles showed that after adjusting for indication bias, there was not enough evidence to suggest a difference in the odds of live birth following blastocyst versus cleavage-stage embryo transfer in the first complete cycle (Cameron et al., 2020), although the majority of the cycles included were performed following culture in atmospheric oxygen, which is known to negatively impact blastocyst outcomes (Gardner, 2016). Although the cumulative live birth rate appears to be similar, blastocyst transfer is associated with a shorter time to pregnancy and to birth and lower cumulative pregnancy loss rates, but also higher transfer cancellation rates compared to cleavage-stage transfer (De Vos et al., 2016; Cornelisse et al., 2024).

The implementation of national strategies towards elective single embryo transfer to decrease multiple birth rates has resulted in increasing use of extended embryo culture (ESHRE Campus Course Report, 2001; Practice Committee of Society for Assisted Reproductive Technology; Practice Committee of American Society for Reproductive Medicine, 2012; Knez et al., 2013; Harbottle et al., 2015; De Geyter et al., 2020; Fouks and Yogev, 2022; ESHRE guideline group on the number of embryos to transfer, 2024).

The development of TE biopsy for PGT has also contributed to the increasing use of blastocyst culture (ESHRE PGT Consortium

Table 9. Consensus scoring system for blastocysts.

	Stage	Description
Stage of expansion	1	Early blastocyst: blastocoel less than half of the volume of the embryo.
	2	Blastocyst: blastocoel that is half of or greater than half of the volume of the embryo.
	3	Full blastocyst: blastocoel completely fills the embryo.
	4	Expanded blastocyst: blastocoel larger than that of the early embryo, with a clearly thinning zona.
	5	Hatching blastocyst: trophectoderm starting to herniate through the zona.
	6	Hatched blastocyst: blastocyst has completely escaped from the zona
	Grade	Description
ICM	A	Prominent, easily discernible, with many cells that are compacted and tightly adhered together.
	B	Easily discernible, with several cells that are loosely grouped together.
	C	Very few cells visible.
	D	No visible cells, or presence of degenerating cells.
TE	A	Many cells forming a cohesive epithelium.
	B	Moderate number of cells forming a loose epithelium.
	C	Few and larger cells with poor epithelia formation.
	D	Sparse or degenerating cells surrounding the ICM

and SIG-Embryology Biopsy Working Group, 2020). Cleavage stage biopsy has been shown to have a negative impact on embryo developmental competence, especially when two blastomeres are removed (Scott et al., 2013). Blastocyst biopsy seems to be safer compared with Day-3 embryo biopsy, as some studies have suggested that removing a small number of TE cells does not affect the embryo implantation or foetal development (Van de Velde et al., 2000; Scott et al., 2013; Tiegs et al., 2021; Cimadomo et al., 2023).

The increasing use of TLT in IVF laboratories, reported in more than 50% of all ART centres responding to our survey (Supplementary Data SII, Fig. 6A), also means that patients are increasingly offered continuous and detailed monitoring of embryo development to blastocyst stage.

Initial concerns about extended embryo culture due to the possible prolonged influence of environmental factors on embryonic epigenetics have appeared to subside (White et al., 2015; Ghosh et al., 2017; Ji et al., 2018), although follow-up studies of children conceived after ART suggest that a possible influence of culture media, culture duration and other laboratory factors on infant health cannot be excluded (Berntsen et al., 2019). Some studies have reported a significantly higher rate of preterm birth after blastocyst transfer compared to cleavage stage transfer (Martins et al., 2016; Wang et al., 2017; Alviggi et al., 2018; Castillo et al., 2020; Cornelisse et al., 2024) while others have reported similar rates (Marconi et al., 2023). In addition, blastocyst transfer appears to be associated with similar or lower risk of small for gestational age (Martins et al., 2016; Raja et al., 2023) and with similar (Siristatidis et al., 2023) or lower congenital anomalies (Raja et al., 2023) compared to cleavage-stage transfer.

One remaining question is whether in poor responders with low zygote numbers, embryo transfer should be done on Day 2, Day 3 or Day 5/6. A retrospective study showed that transferring embryos on Day 2 versus Day 3 in this patient group does not affect early pregnancy outcomes and suggested the flexibility in scheduling the day of transfer at the convenience of both the patient and the centre (Sacha et al., 2018). According to another study, there is no difference in clinical pregnancy rates after fresh Day-3 or Day-5 embryo transfer in patients with 5 or fewer zygotes (Dirican et al., 2022). However, those with six or more zygotes can benefit from blastocyst transfer due to better selection options. Larger prospective studies on live birth rates also taking into account maternal age are needed to provide a conclusive answer to the above question.

Technical considerations for extended embryo culture

The success of extended embryo culture relies on crucial parameters, such as reduced oxygen concentration, optimal pH, temperature and osmolality (Gardner and Lane, 1997). Blastocyst culture affects logistics and workflow, as well as technical requirements in the laboratory, such as incubator type and capacity, frequency of embryo assessment, and—if performed—annotation of morphokinetics and culture media renewal. Success also depends on stable culture conditions and an efficient blastocyst vitrification programme (Swain, 2019; Cairo Consensus Group, 2020). Therefore, the ART centre's capacity to ensure appropriate conditions for blastocyst culture should be proven. A blastocyst culture approach should be introduced starting first with good responder patients and, after appropriate blastocyst development rate and clinical outcomes are obtained, gradual wider applications can be offered to other groups of IVF patients (Gardner and Lane, 2018; De Croo et al., 2020). The success of the blastocyst vitrification programme should be self-verified by the IVF laboratory by tracking key performance indicators. The reference rates for blastocyst cryosurvival are expected to be $\geq 90\%$ for competency and $\geq 99\%$ for benchmark (ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine, 2017). Due to greater experience with blastocyst vitrification, the rate of degeneration during warming is now lower than that estimated in a previous cryopreservation consensus (Alpha Scientists In Reproductive Medicine, 2012).

Modern benchtop incubators with individual chambers represent a safer incubator design and provide a faster recovery time of all physico-chemical parameters after door openings compared to older 'big-box' incubators (Kovačić, 2021). However, in the case of prolonged and continuous culture of embryos, possible changes in osmolality and pH over time must also be monitored (Swain, 2019), also taking into consideration the type of dishware, culture drop size and oil overlay.

Incubators with integrated TLT allow continuous observation of the morphokinetics of developing embryos with uninterrupted incubation throughout the preimplantation period (Meseguer et al., 2012). A good practice recommendation paper including a systematic assessment of how to approach and introduce TLT for IVF was published to provide centres with technical advice (ESHRE Working group on Time-lapse technology et al., 2020).

Due to the overwhelming evidence of the detrimental effect of atmospheric oxygen concentration on embryonic development (Gardner, 2016), the use of reduced oxygen is now considered standard practice, especially for extended incubation of

embryos to blastocyst stage (Kovačić, 2012; De los Santos et al., 2016).

Frequency of embryo assessment: rationale

While the accuracy of assessing blastomere cleavages is important, laboratories with limited number of incubators should carefully consider certain limitations and prioritize the safety and quality of the embryo culture conditions. More frequent opening of incubators may have a negative impact on embryonic development (Gardner and Lane, 1996; Zhang et al., 2010; Nguyen et al., 2018). In such situations, assessing morphology only at the end of the culture period may be considered, with no or few intermediate checks on their development.

If it is decided to practice short-term embryo culture in IVF cycles with large numbers of zygotes, then a more detailed and frequent assessment of embryo morphology might improve selection of embryos by the ranking scheme given in this paper.

Consensus points

- Extended embryo culture is an accepted and standard practice.
- The duration of embryo culture and frequency of static embryo observations must be adjusted to the equipment in the laboratory and staff skill, ensuring minimal changes in culture conditions that could affect embryo development.

Conclusion

This consensus paper provides updated recommendations on criteria and terminology for assessing oocyte, zygote, cleavage-stage embryo, morula and blastocyst development, based on a thorough review of evidence accumulated over the past decade. Critical information gained from application of TLT has provided the impetus for revised timings of developmental milestones,

Table 10. List of recommendations.

RECOMMENDATIONS	
OOCYTE ASSESSMENT	<ul style="list-style-type: none"> • Giant oocytes should be excluded from clinical use. • The use of small/large oocytes and IVM-rescued oocytes should be documented for prognostic and traceability purposes due to their apparently lower developmental potential. • Finally, embryos derived from MII oocytes free of large or multiple vacuoles, SER-a, and very large first PB should be prioritized for clinical use. • Prenatal follow-up and the follow-up of babies born from oocytes with atypical phenotypes and rescue IVM demands attention.
ZYGOTE STATE ASSESSMENT	<ul style="list-style-type: none"> • Assessment of PN number should be carried out between 16 and 17 hpi in both conventional IVF and ICSI cases. • Zygotes with 2PN should be prioritized for clinical use. • 2.1PN and 1PN zygotes from IVF or ICSI may be considered for clinical use with appropriate counselling, especially if associated with PGT-A technology appropriate for biparental diploidy assessment. • The clinical use of 3PN zygotes is not recommended, while pre-clinical or pilot clinical studies should be encouraged. • Dynamic features such as PN size, PN position and juxtaposition, NPB pattern, and cytoplasmic halo cannot be accurately assessed during static observations. Thus, they cannot be consistently used as biomarkers of viability.
DAY -1, -2 & -3 EMBRYO ASSESSMENT	<ul style="list-style-type: none"> • 2-cell embryos on Day-1, 4-cell embryos on Day- 2, 8-cell embryos on Day-3 showing <10% fragmentation, mononucleation, and stage-specific cell size should be prioritized in case of cleavage stage embryo transfer or cryopreservation. • Cleavage stage embryos with atypical features such as extensive fragmentation, multinucleation, vacuoles, cytoplasmic granularity, membrane, and zona irregularities, can be considered suitable for clinical use. However, extended culture of these embryos for further evaluation should be considered.
DAY-4 EMBRYO ASSESSMENT	<ul style="list-style-type: none"> • Day-4 embryos showing full compaction or early cavitation should be prioritized in case of Day-4 transfer or vitrification. • Embryos with partial compaction can form blastocysts and should be considered for clinical use. Extended culture of these embryos for further evaluation should be considered.
DAY-5, -6 & -7 EMBRYO ASSESSMENT	<ul style="list-style-type: none"> • The Gardner grading system for blastocyst scoring (Table 8) should be used. This system is distinguished from the prior Consensus grading by using letters for the ICM/TE grades and adding additional expansion stages (e.g. hatched blastocyst). • Non-viable blastocysts should be graded as "D" as opposed to "C" based on degenerative features or absence of a distinct ICM. • The common features that are clearly associated with implantation potential include day of blastocyst formation (Day 4-7), stage of expansion (3,4,5,6), and grade of ICM (A, B, C) and TE (A, B, C). • Blastocysts with grade C ICM and/or TE and Day 7 blastocysts can be viable and could be considered suitable for clinical use. • Blastocysts with 2 ICM indicating potential monozygotic twinning should not be transferred without thorough patient counselling. • Assigning relative importance of each variable requires systematic multivariate analysis with a large dataset and is further complicated when assessing fresh versus frozen untested and euploid blastocysts.
DURATION OF EMBRYO CULTURE AND FREQUENCY OF ASSESSMENTS	<ul style="list-style-type: none"> • Extended embryo culture is an accepted and standard practice. • The length of embryo culture and frequency of static embryo observations must be adjusted to the equipment in the laboratory and staff skill, ensuring minimal changes in culture conditions that could affect embryo development.

Table 11. List of knowledge gaps and recommendations for future research.

	Knowledge gap	Recommendations for future research
Expected timeline	It is unknown how and whether artificial intelligence-based analyses and selection algorithms will evolve or deal with data heterogeneity.	Develop more advanced analytic tools to provide the facility to identify the most viable embryo(s) from a cohort and an estimation of the likelihood of each embryo leading to live birth.
Oocyte assessment	The body of evidence to date is based almost exclusively on qualitative (presence/absence) analyses and exclude an objective description of each dysmorphism. The impact of different ovarian stimulation protocols/responses on oocyte parameters has not been fully evaluated.	Future standardized and quantitative analyses should be conducted on oocyte morphology, thereby filling important gaps in knowledge. Further studies using artificial intelligence for oocyte assessment might be useful.
Zygote stage assessment	TLT has highlighted complex changes over time of the majority of relevant morphokinetic parameters, such as PN size and position, NPB patterning and cytoplasmic halo. Use of such parameters to predict embryo developmental competence remains elusive, probably because morphokinetic abnormalities occurring at fertilisation may be compensated by the outstanding developmental plasticity of the human embryo.	Use of TLT and allied technologies, namely image analysis and artificial intelligence to decrypt the developmental significance of fertilisation biomarkers, such as NPB patterning. This is expected to lead to novel criteria for embryo ranking and perhaps for the prediction of blastocyst aneuploidy. Use of TLT and PGT-A to distinguish haploid/triploid from diploid 1PN and 3PN zygotes, thereby identifying potentially viable embryos that would be otherwise discarded.
Day -1, -2 & -3 embryo assessment	Insight into what may be considered optimal timing for cleavage stage embryo evaluation is still lacking. Questions surrounding the significance of multinucleation, the number of nuclei and the number of affected cells and the developmental stage when this condition appears remain largely unanswered. There is a crucial gap in knowledge concerning the criteria for exclusion of embryos from selection for clinical use.	Further studies using TLT are expected to provide a deeper understanding of the association between time of assessment, morphological features, and clinical outcomes.
Day-4 embryo assessment	It is currently unknown whether and to what extent type and composition of culture media (e.g. Ca ²⁺ , Mg ²⁺) might influence compaction timing and phenotypes. Little information is available on premature compaction behaviour as early as the 2- to 4-cell stages.	Explore the underlying cellular mechanisms that can explain compaction timing and blastomere exclusion/extrusion processes.
Day-5, -6 & -7 embryo assessment	A best practice for establishing a clinic-specific ranking of blastocysts based on morphology and time of development and in-house validation of established algorithms before use is lacking.	Develop objective measures of blastocyst quality to improve the accuracy of blastocyst scoring and ranking, though early reports have not shown an improvement with either of these methodologies. Identify markers of viability beyond morphology and bright-field microscopy to improve non-invasive blastocyst assessment.
Duration of embryo culture and frequency of assessments	Evidence is lacking on the effectiveness of non-selective use of extended embryo culture in all patients.	Assess whether more frequent observations of embryos during prolonged culture improves embryo selection or clinical efficacy of the procedure.

greater consideration of the influence of insemination methods on early embryogenesis, and presentation of a broader spectrum of atypical morphology detected with time-lapse imaging. The collated recommendations (Table 10) aim to promote standardized embryo evaluation practices to better predict viability and optimize embryo ranking and selection for clinical use. Notwithstanding the progress of the past decade, several knowledge gaps remain (Table 11) concerning the clinical value of specific morphological and morphokinetic parameters that warrant further investigation and scrutiny. Undoubtedly, the next decade will bring a more substantial incorporation of AI in the ART laboratory, offering solutions to the perpetually challenging problem of viable gamete and embryo selection.

Lastly, by combining expertise and experience across institutions and geographical regions, international collaborative efforts such as that represented by this consensus paper can contribute to improving research consistency, clinical practice, and most

importantly, outcomes for patients seeking assisted reproduction.

Supplementary data

Supplementary data are available at *Human Reproduction* online.

Data availability

This article conducts a literature review of existing research records, and no new data were generated or analysed in support of this manuscript.

Acknowledgements

The authors would like to thank Katie Best (Phastar, London) for her assistance in statistical analysis of observation timings for

the different developmental stages of the embryo represented in Table 1 of this manuscript, and Alessandro Bartolacci (IRCCS San Raffaele Scientific institute, Milan) and Marilena Taggi (IVI-RMA Global Research Alliance, Rome) for their assistance in the literature search for oocyte assessment. They would also like to thank all the ALPHA and ESHRE members who have answered the survey, and those who contributed to the stakeholder review.

Authors' role

This document resulted from a joint effort of ESHRE and ALPHA. M.A. and G.C. co-chaired the working group and contributed equally to writing the paper and critical reading. G.A., A.C., T.E., D.K.G., B.K., K.L., D.M., L.R., and I.S., listed in alphabetical order, as working group members, contributed equally to writing the paper and critical reading. A.A., B.B., M.J.D.S., and C.M. were invited experts to the consensus meeting and contributed equally to reviewing the paper and critical reading. S.M. and N.V., as methodological experts, performed data collection and analysis for the survey, literature searches, provided methodological support and coordinated the development of this manuscript.

Funding

The consensus meeting and writing of the paper was supported by funds from ESHRE and ALPHA.

Conflict of interest

G.C. declared payments or honoraria for lectures from Gedeon Richter and Cooper Surgical. A.C. declared text book royalties (Mastering Clinical Embryology, published 2024), consulting fees from Cooper Surgical, Gedeon Richter and TMRW Life Sciences, honoraria for lectures from Merck, Ferring, and Gedeon Richter, and participation in the HFEA Scientific Advances Committee; she also disclosed being treasurer and vice-president of Alpha Scientists in Reproductive Medicine, a shareholder in Care Fertility Limited and Fertile Mind Limited, and having stock options in TMRW Life Sciences and U-Ploid Biotechnology Ltd. L. R. declared consulting fees from Organon, payments or honoraria for lectures from Merck, Organon, IBSA, Finox, Geden Richter, Origio, Organon, Ferring, Foundation IVI; she also disclosed being a member of the Advisory Scientific Board of IVIRMA (Paid) and a member of the Advisory Scientific Board of Nterilizer (unpaid). I. S. declared payments or honoraria for lectures from Vitrolife and Cooper Surgical, and stock options from Alife Health. M.A. declared payments or honoraria for lectures from Vitrolife and support for attending meetings from Vitrolife and Cooper Surgical (both unrelated to this manuscript). The other authors have no conflicts of interest to declare.

References

Abdala A, Elkhatib I, Bayram A, Aranz A, El-Damen A, Melado L, Lawrenz B, Fatemi HM, De Munck N. Day 5 vs day 6 single euploid blastocyst frozen embryo transfers: which variables do have an impact on the clinical pregnancy rates? *J Assist Reprod Genet* 2022; **39**:379–388.

Abdala A, Elkhatib I, Bayram A, El-Damen A, Melado L, Nogueira D, Lawrenz B, Fatemi HM. Reproductive outcomes with delayed blastocyst development: the clinical value of day 7 euploid blastocysts in frozen embryo transfer cycles. *Zygote* 2023; **31**:588–595.

Aguilar J, Motato Y, Escribá MJ, Ojeda M, Muñoz E, Meseguer M. The human first cell cycle: impact on implantation. *Reprod Biomed Online* 2014; **28**:475–484.

Aguilar J, Rubio I, Muñoz E, Pellicer A, Meseguer M. Study of nucleation status in the second cell cycle of human embryo and its impact on implantation rate. *Fertil Steril* 2016; **106**:291–299.e2.

Ahlström A, Berntsen J, Johansen M, Bergh C, Cimadomo D, Hardarson T, Lundin K. Correlations between a deep learning-based algorithm for embryo evaluation with cleavage-stage cell numbers and fragmentation. *Reprod Biomed Online* 2023; **47**:103408.

Ahlström A, Lundin K, Lind AK, Gunnarsson K, Westlander G, Park H, Thurin-Kjellberg A, Thorsteinsdóttir SA, Einarsson S, Åström M et al A double-blind randomized controlled trial investigating a time-lapse algorithm for selecting Day 5 blastocysts for transfer. *Hum Reprod* 2022; **37**:708–717.

Ahlström A, Park H, Bergh C, Selleskog U, Lundin K. Conventional morphology performs better than morphokinetics for prediction of live birth after day 2 transfer. *Reprod Biomed Online* 2016; **33**:61–70.

Ahlström A, Westin C, Reisner E, Wikland M, Hardarson T. Trophoctoderm morphology: an important parameter for predicting live birth after single blastocyst transfer. *Hum Reprod* 2011; **26**:3289–3296.

Ahlström A, Westin C, Wikland M, Hardarson T. Prediction of live birth in frozen-thawed single blastocyst transfer cycles by pre-freeze and post-thaw morphology. *Hum Reprod* 2013; **28**:1199–1209.

Akarsu C, Çağlar G, Vicdan K, Sözen E, Biberoglu K. Smooth endoplasmic reticulum aggregations in all retrieved oocytes causing recurrent multiple anomalies: case report. *Fertil Steril* 2009; **92**:1496.e1–1496.e3.

Alikani M. Monozygotic twinning following assisted conception: an analysis of 81 consecutive cases. *Hum Reprod* 2003; **18**:1937–1943.

Alikani M. Epithelial cadherin distribution in abnormal human preimplantation embryos. *Hum Reprod* 2005; **20**:3369–3375.

Alikani M, Calderon G, Tomkin G, Garrisi J, Kokot M, Cohen J. Cleavage anomalies in early human embryos and survival after prolonged culture in-vitro. *Hum Reprod* 2000; **15**:2634–2643.

Alikani M, Cohen J, Tomkin G, Garrisi GJ, Mack C, Scott RT. Human embryo fragmentation in vitro and its implications for pregnancy and implantation. *Fertil Steril* 1999; **71**:836–842.

Alikani M, Palermo G, Adler A, Bertoli M, Blake M, Cohen J. Intracytoplasmic sperm injection in dysmorphic human oocytes. *Zygote* 1995; **3**:283–288.

Almeida PA, Bolton VN. The relationship between chromosomal abnormality in the human preimplantation embryo and development in vitro. *Reprod Fertil Dev* 1996; **8**:235–241.

Alpha Scientists In Reproductive Medicine. The Alpha consensus meeting on cryopreservation key performance indicators and benchmarks: proceedings of an expert meeting. *Reproductive Biomedicine Online* 2012; **25**:146–167.

Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group Embryology. Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Reprod Biomed Online* 2011; **22**:632–646.

Alvigi C, Conforti A, Carbone IF, Borrelli R, de Placido G, Guerriero S. Influence of cryopreservation on perinatal outcome after blastocyst- vs cleavage-stage embryo transfer: systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2018; **51**:54–63.

Anagnostopoulou C, Maldonado Rosas I, Singh N, Gugnani N, Chockalingham A, Singh K, Desai D, Darbandi M, Manoharan M, Darbandi S et al Oocyte quality and embryo selection strategies:

- a review for the embryologists, by the embryologists. *Panminerva Med* 2022;**64**:171–184.
- Antczak M, Van Blerkom J. Temporal and spatial aspects of fragmentation in early human embryos: possible effects on developmental competence and association with the differential elimination of regulatory proteins from polarized domains. *Hum Reprod* 1999;**14**:429–447.
- Araki E, Itoi F, Honnma H, Asano Y, Oguri H, Nishikawa K. Correlation between the pronucleus size and the potential for human single pronucleus zygotes to develop into blastocysts: 1PN zygotes with large pronuclei can expect an embryo development to the blastocyst stage that is similar to the development of 2PN zygotes. *J Assist Reprod Genet* 2018;**35**:817–823.
- Arce JC, Ziebe S, Lundin K, Janssens R, Helmgaard L, Sørensen P. Interobserver agreement and intraobserver reproducibility of embryo quality assessments. *Hum Reprod* 2006;**21**:2141–2148.
- Armstrong S, Bhide P, Jordan V, Pacey A, Marjoribanks J, Farquhar C. Time-lapse systems for embryo incubation and assessment in assisted reproduction. *Cochrane Database Syst Rev* 2019;**5**:Cd011320.
- Arroyo G, Santaló J, Boada M, Parriego M, Rodríguez I, Coroleu B, Barri PN, Veiga A. Does early cleavage correlate with chromosome constitution in human preimplantation embryos? *Med Reprod Embriol Clín* 2015;**2**:31–39.
- Asami M, Lam BYH, Ma MK, Rainbow K, Braun S, VerMilyea MD, Yeo GSH, Perry ACF. Human embryonic genome activation initiates at the one-cell stage. *Cell Stem Cell* 2022;**29**:209–216.
- Ashrafi M, Karimian L, Eftekhari-Yazdi P, Hasani F, Arabipoor A, Bahmanabadi A, Akhond MR. Effect of oocyte dysmorphisms on intracytoplasmic sperm injection cycle outcomes in normal ovarian responders. *J Obstet Gynaecol Res* 2015;**41**:1912–1920.
- Aslan Öztürk S, Cincik M, Donmez Cakil Y, Sayan S, Selam B. Early compaction might be a parameter to determine good quality embryos and day of embryo transfer in patients undergoing intracytoplasmic sperm injection. *Cureus* 2022;**14**:e23593.
- Awadalla MS, Agarwal R, Ho JR, McGinnis LK, Ahmady A. Effect of trophectoderm biopsy for PGT-A on live birth rate per embryo in good prognosis patients. *Arch Gynecol Obstet* 2022a;**306**:1321–1327.
- Awadalla MS, Ho JR, McGinnis LK, Ahmady A, Cortessis VK, Paulson RJ. Embryo morphology and live birth in the United States. *FS Rep* 2022b;**3**:131–137.
- Azzarello A, Hoest T, Mikkelsen AL. The impact of pronuclei morphology and dynamicity on live birth outcome after time-lapse culture. *Hum Reprod* 2012;**27**:2649–2657.
- Bakkensen JB, Brady P, Carusi D, Romanski P, Thomas AM, Racowsky C. Association between blastocyst morphology and pregnancy and perinatal outcomes following fresh and cryopreserved embryo transfer. *J Assist Reprod Genet* 2019;**36**:2315–2324.
- Balaban B, Ata B, Isiklar A, Yakin K, Urman B. Severe cytoplasmic abnormalities of the oocyte decrease cryosurvival and subsequent embryonic development of cryopreserved embryos. *Hum Reprod* 2008;**23**:1778–1785.
- Balaban B, Urman B. Effect of oocyte morphology on embryo development and implantation. *Reprod Biomed Online* 2006;**12**:608–615.
- Balaban B, Urman B, Sertac A, Alatas C, Aksoy S, Mercan R. Oocyte morphology does not affect fertilization rate, embryo quality and implantation rate after intracytoplasmic sperm injection. *Hum Reprod* 1998;**13**:3431–3433.
- Balakier H, Bouman D, Sojecki A, Librach C, Squire JA. Morphological and cytogenetic analysis of human giant oocytes and giant embryos. *Hum Reprod* 2002;**17**:2394–2401.
- Balakier H, Cadesky K. The frequency and developmental capability of human embryos containing multinucleated blastomeres. *Hum Reprod* 1997;**12**:800–804.
- Balakier H, Sojecki A, Motamedi G, Librach C. Time-dependent capability of human oocytes for activation and pronuclear formation during metaphase II arrest. *Hum Reprod* 2004;**19**:982–987.
- Balakier H, Sojecki A, Motamedi G, Librach C. Impact of multinucleated blastomeres on embryo developmental competence, morphokinetics, and aneuploidy. *Fertil Steril* 2016;**106**:608–614.e2.
- Bamford T, Barrie A, Montgomery S, Dhillon-Smith R, Campbell A, Easter C, Coomarasamy A. Morphological and morphokinetic associations with aneuploidy: a systematic review and meta-analysis. *Hum Reprod Update* 2022;**28**:656–686.
- Bamford T, Easter C, Montgomery S, Smith R, Dhillon-Smith RK, Barrie A, Campbell A, Coomarasamy A. A comparison of 12 machine learning models developed to predict ploidy, using a morphokinetic meta-dataset of 8147 embryos. *Hum Reprod* 2023;**38**:569–581.
- Barberet J, Bruno C, Valot E, Antunes-Nunes C, Jonval L, Chammas J, Choux C, Ginod P, Sagot P, Soudry-Faure A et al Can novel early non-invasive biomarkers of embryo quality be identified with time-lapse imaging to predict live birth? *Hum Reprod* 2019;**34**:1439–1449.
- Barnes J, Brendel M, Gao VR, Rajendran S, Kim J, Li Q, Malmsten JE, Sierra JT, Zisimopoulos P, Sigaras A et al A non-invasive artificial intelligence approach for the prediction of human blastocyst ploidy: a retrospective model development and validation study. *Lancet Digit Health* 2023;**5**:e28–e40.
- Barrie A, Homburg R, McDowell G, Brown J, Kingsland C, Troup S. Examining the efficacy of six published time-lapse imaging embryo selection algorithms to predict implantation to demonstrate the need for the development of specific, in-house morphokinetic selection algorithms. *Fertil Steril* 2017a;**107**:613–621.
- Barrie A, Homburg R, McDowell G, Brown J, Kingsland C, Troup S. Preliminary investigation of the prevalence and implantation potential of abnormal embryonic phenotypes assessed using time-lapse imaging. *Reprod Biomed Online* 2017b;**34**:455–462.
- Barrie A, McDowell G, Troup S. An investigation into the effect of potential confounding patient and treatment parameters on human embryo morphokinetics. *Fertil Steril* 2021a;**115**:1014–1022.
- Barrie A, Smith R, Campbell A, Fishel S. Optimisation of the timing of fertilisation assessment for oocytes cultured in standard incubation: lessons learnt from time-lapse imaging of 78 348 embryos. *Hum Reprod* 2021b;**36**:2840–2847.
- Bartolacci A, Intra G, Cotichio G, dell'Aquila M, Patria G, Borini A. Does morphological assessment predict oocyte developmental competence? A systematic review and proposed score. *J Assist Reprod Genet* 2022;**39**:3–17.
- Bassil R, Casper RF, Meriano J, Smith R, Haas J, Mehta C, Orvieto R, Zilberberg E. Can oocyte diameter predict embryo quality? *Reprod Sci* 2021;**28**:904–908.
- Baxter Bendus AE, Mayer JF, Shipley SK, Catherino WH. Interobserver and intraobserver variation in day 3 embryo grading. *Fertil Steril* 2006;**86**:1608–1615.
- Bellver J. BMI and miscarriage after IVF. *Curr Opin Obstet Gynecol* 2022;**34**:114–121.
- Berntsen J, Rimestad J, Lassen JT, Tran D, Kragh MF. Robust and generalizable embryo selection based on artificial intelligence and time-lapse image sequences. *PLoS One* 2022;**17**:e0262661.
- Berntsen S, Söderström-Anttila V, Wennerholm UB, Laivuori H, Loft A, Oldereid NB, Romundstad LB, Bergh C, Pinborg A. The health of children conceived by ART: ‘the chicken or the egg?’. *Hum Reprod Update* 2019;**25**:137–158.

- Bertrand E, Van den Bergh M, Englert Y. Does zona pellucida thickness influence the fertilization rate? *Hum Reprod* 1995; **10**:1189–1193.
- Bickendorf K, Qi F, Peirce K, Natalwala J, Chapple V, Liu Y. Spontaneous collapse as a prognostic marker for human blastocysts: a systematic review and meta-analysis. *Hum Reprod* 2023; **38**:1891–1900.
- Bodri D, Kato R, Kondo M, Hosomi N, Katsumata Y, Kawachiya S, Matsumoto T. Time-lapse monitoring of zona pellucida-free embryos obtained through in vitro fertilization: a retrospective case series. *Fertil Steril* 2015; **103**:e35.
- Bodri D, Kawachiya S, Sugimoto T, Yao Serna J, Kato R, Matsumoto T. Time-lapse variables and embryo gender: a retrospective analysis of 81 live births obtained following minimal stimulation and single embryo transfer. *J Assist Reprod Genet* 2016; **33**:589–596.
- Bormann CL, Thirumalaraju P, Kanakasabapathy MK, Kandula H, Souter I, Dimitriadis I, Gupta R, Pooniwala R, Shafiee H. Consistency and objectivity of automated embryo assessments using deep neural networks. *Fertil Steril* 2020; **113**:781–787.e1.
- Boucret L, Tramon LÉA, Riou J, Ferré-L'Hôtelier V, Bouet P-E, May-Panloup P. Influence of diminished ovarian reserve on early embryo morphokinetics during in vitro fertilization: a time-lapse study. *JCM* 2022; **11**:7173.
- Bourdon M, Pocate-Cheriet K, Finet de Bantel A, Grzegorzczuk-Martin V, Amar Hoffet A, Arbo E, Poulain M, Santulli P. Day 5 versus Day 6 blastocyst transfers: a systematic review and meta-analysis of clinical outcomes. *Hum Reprod* 2019; **34**:1948–1964.
- Bradley CK, Traversa MV, Hobson N, Gee AJ, McArthur SJ. Clinical use of monopronucleated zygotes following blastocyst culture and preimplantation genetic screening, including verification of biparental chromosome inheritance. *Reprod Biomed Online* 2017; **34**:567–574.
- Braga DP, Halpern G, Setti AS, Figueira RC, Iaconelli A Jr, Borges E Jr. The impact of food intake and social habits on embryo quality and the likelihood of blastocyst formation. *Reprod Biomed Online* 2015; **31**:30–38.
- Braga DPAF, Setti AS, Figueira R, Iaconelli A, Borges E Jr. The combination of pronuclear and blastocyst morphology: a strong prognostic tool for implantation potential. *J Assist Reprod Genet* 2013; **30**:1327–1332.
- Braude P, Bolton V, Moore S. Human gene expression first occurs between the four- and eight-cell stages of preimplantation development. *Nature* 1988; **332**:459–461.
- Bredbacka P, Capalbo A, Kananen K, Picchetta L, Tomás C. Healthy live birth following embryo transfer of a blastocyst of tetrapronuclear (4PN) origin: a case report. *Hum Reprod* 2023; **38**:1700–1704.
- Busnelli A, Dallagiovanna C, Reschini M, Paffoni A, Fedele L, Somigliana E. Risk factors for monozygotic twinning after in vitro fertilization: a systematic review and meta-analysis. *Fertil Steril* 2019; **111**:302–317.
- Cameron NJ, Bhattacharya S, McLernon DJ. Cumulative live birth rates following blastocyst- versus cleavage-stage embryo transfer in the first complete cycle of IVF: a population-based retrospective cohort study. *Hum Reprod* 2020; **35**:2365–2374.
- Campbell A, Cohen J, Ivani K, Morbeck D, Palmer G, Mortimer S. The in vitro fertilization laboratory: teamwork and teaming. *Fertil Steril* 2022a; **117**:27–32.
- Campbell A, Fishel S, Bowman N, Duffy S, Sedler M, Hickman CF. Modelling a risk classification of aneuploidy in human embryos using non-invasive morphokinetics. *Reprod Biomed Online* 2013; **26**:477–485.
- Campbell AJ, Petersen Dr BM, Smith R, Barrie A. Prediction of blastulation, embryo utilisation and live birth from single morphological or morphokinetic variables: analysis of 31,323 embryos gives insights for selection and algorithm development. *Fertil Steril* 2022b; **118**:e138.
- Capalbo A, Rienzi L, Cimadomo D, Maggiulli R, Elliott T, Wright G, Nagy ZP, Ubaldi FM. Correlation between standard blastocyst morphology, euploidy and implantation: an observational study in two centers involving 956 screened blastocysts. *Hum Reprod* 2014; **29**:1173–1181.
- Capalbo A, Treff N, Cimadomo D, Tao X, Ferrero S, Vaiarelli A, Colamaria S, Maggiulli R, Orlando G, Scarica C et al Abnormally fertilized oocytes can result in healthy live births: improved genetic technologies for preimplantation genetic testing can be used to rescue viable embryos in in vitro fertilization cycles. *Fertil Steril* 2017; **108**:1007–1015.e3.
- Castillo CM, Johnstone ED, Horne G, Falconer DA, Troup SA, Cutting R, Sharma V, Brison DR, Roberts SA. Associations of IVF singleton birthweight and gestation with clinical treatment and laboratory factors: a multicentre cohort study. *Hum Reprod* 2020; **35**:2860–2870.
- Cauffman G, Verheyen G, Tournaye H, Van de Velde H. Developmental capacity and pregnancy rate of tetrahedral- versus non-tetrahedral-shaped 4-cell stage human embryos. *J Assist Reprod Genet* 2014; **31**:427–434.
- Cavazza T, Takeda Y, Politi AZ, Aushev M, Aldag P, Baker C, Choudhary M, Bucevičius J, Lukinavičius G, Elder K et al Parental genome unification is highly error-prone in mammalian embryos. *Cell* 2021; **184**:2860–2877.e22.
- Cecchele A, Cermisoni GC, Giacomini E, Pinna M, Vigano P. Cellular and molecular nature of fragmentation of human embryos. *IJMS* 2022; **23**:1349.
- Chamayou S, Alecci C, Ragolia C, Storaci G, Maglia E, Russo E, Guglielmino A. Comparison of in-vitro outcomes from cryopreserved oocytes and sibling fresh oocytes. *Reprod Biomed Online* 2006; **12**:730–736.
- Chatzimeletiou K, Morrison EE, Prapas N, Prapas Y, Handyside AH. Spindle abnormalities in normally developing and arrested human preimplantation embryos in vitro identified by confocal laser scanning microscopy. *Hum Reprod* 2005; **20**:672–682.
- Chavez-Badiola A, Flores-Saiffe-Farías A, Mendizabal-Ruiz G, Drakeley AJ, Cohen J. Embryo Ranking Intelligent Classification Algorithm (ERICA): artificial intelligence clinical assistant predicting embryo ploidy and implantation. *Reprod Biomed Online* 2020; **41**:585–593.
- Chavez SL, Loewke KE, Han J, Moussavi F, Colls P, Munne S, Behr B, Reijo Pera RA. Dynamic blastomere behaviour reflects human embryo ploidy by the four-cell stage. *Nat Commun* 2012; **3**:1251.
- Chen X, Zhang J, Wu X, Cao S, Zhou L, Wang Y, Chen X, Lu J, Zhao C, Chen M et al Trophectoderm morphology predicts outcomes of pregnancy in vitrified-warmed single-blastocyst transfer cycle in a Chinese population. *J Assist Reprod Genet* 2014; **31**:1475–1481.
- Chen ZQ, Wang Y, Ng EHY, Zhao M, Pan JP, Wu HX, Teng XM. A randomized triple blind controlled trial comparing the live birth rate of IVF following brief incubation versus standard incubation of gametes. *Hum Reprod* 2019; **34**:100–108.
- Cimadomo D, Capalbo A, Scarica C, Sosa Fernandez L, Rienzi L, Ciriminna R, Minasi MG, Novelli A, De Santis L, Zuccarello D. When embryology meets genetics: the definition of developmentally incompetent preimplantation embryos (DIPE)-the consensus of two Italian scientific societies. *J Assist Reprod Genet* 2021; **38**:319–331.
- Cimadomo D, Marconetto A, Trio S, Chiappetta V, Innocenti F, Albricci L, Erlich I, Ben-Meir A, Har-Vardi I, Kantor B et al Human blastocyst spontaneous collapse is associated with worse morphological quality and higher degeneration and aneuploidy rates:

- a comprehensive analysis standardized through artificial intelligence. *Hum Reprod* 2022a;**37**:2291–2306.
- Cimadomo D, Rienzi L, Conforti A, Forman E, Canosa S, Innocenti F, Poli M, Hynes J, Gemmell L, Vaiarelli A et al Opening the black box: why do euploid blastocysts fail to implant? A systematic review and meta-analysis. *Hum Reprod Update* 2023;**29**:570–633.
- Cimadomo D, Soscia D, Casciani V, Innocenti F, Trio S, Chiappetta V, Albricci L, Maggiulli R, Erlich I, Ben-Meir A et al How slow is too slow? A comprehensive portrait of Day 7 blastocysts and their clinical value standardized through artificial intelligence. *Hum Reprod* 2022b;**37**:1134–1147.
- Ciotti PM, Notarangelo L, Morselli-Labate AM, Felletti V, Porcu E, Venturoli S. First polar body morphology before ICSI is not related to embryo quality or pregnancy rate. *Hum Reprod* 2004;**19**:2334–2339.
- Ciray HN, Campbell A, Agerholm IE, Aguilar J, Chamayou S, Esbert M, Sayed S; Time-Lapse User Group. Proposed guidelines on the nomenclature and annotation of dynamic human embryo monitoring by a time-lapse user group. *Hum Reprod* 2014;**29**:2650–2660.
- Cohen J, Levron J, Palermo GD, Munné S, Adler A, Alikani M, Schattman G, Sultan K, Willadsen S. Atypical activation and fertilization patterns in humans. *Theriogenology* 1995;**43**:129–140.
- Cornelisse S, Fleischer K, van der Westerlaken L, de Bruin J-P, Vergouw C, Koks C, Derhaag J, Visser J, van Echten-Arends J, Slappendel E et al Cumulative live birth rate of a blastocyst versus cleavage stage embryo transfer policy during in vitro fertilisation in women with a good prognosis: multicentre randomised controlled trial. *BMJ* 2024;**386**:e080133.
- Coticchio G, E Zoe K, Lagalla C, Shimazaki K, Ohata K, Ninomiya M, Wakabayashi N, Okimura T, Uchiyama K, Kato K et al Perturbations of morphogenesis at the compaction stage affect blastocyst implantation and live birth rates. *Hum Reprod* 2021;**36**:918–928.
- Coticchio G, E Zoe K, Lagalla C, Zacà C, Borini A, Kato K. The destinies of human embryos reaching blastocyst stage between Day 4 and Day 7 diverge as early as fertilization. *Hum Reprod* 2023;**38**:1690–1699.
- Coticchio G, Lagalla C, Sturmey R, Pennetta F, Borini A. The enigmatic morula: mechanisms of development, cell fate determination, self-correction and implications for ART. *Hum Reprod Update* 2019;**25**:422–438.
- Coticchio G, Mignini Renzini M, Novara PV, Lain M, De Ponti E, Turchi D, Fadini R, Dal Canto M. Focused time-lapse analysis reveals novel aspects of human fertilization and suggests new parameters of embryo viability. *Hum Reprod* 2018;**33**:23–31.
- Cuevas Saiz I, Carme Pons Gatell M, Vargas MC, Delgado Mendive A, Rives Enedáguila N, Moragas Solanes M, Carrasco Canal B, Teruel López J, Busquets Bonet A, Hurtado de Mendoza Acosta MV. The Embryology Interest Group: updating ASEBIR's morphological scoring system for early embryos, morulae and blastocysts. *Med Reprod Embriol Clín* 2018;**5**:42–54.
- Currie CE, Ford E, Benham Whyte L, Taylor DM, Mihalas BP, Erent M, Marston AL, Hartshorne GM, McAinsh AD. The first mitotic division of human embryos is highly error prone. *Nat Commun* 2022;**13**:6755.
- Dal Canto M, Brambillasca F, Mignini Renzini M, Coticchio G, Merola M, Lain M, De Ponti E, Fadini R. Cumulus cell-oocyte complexes retrieved from antral follicles in IVM cycles: relationship between COCs morphology, gonadotropin priming and clinical outcome. *J Assist Reprod Genet* 2012a;**29**:513–519.
- Dal Canto M, Coticchio G, Mignini Renzini M, De Ponti E, Novara PV, Brambillasca F, Comi R, Fadini R. Cleavage kinetics analysis of human embryos predicts development to blastocyst and implantation. *Reprod Biomed Online* 2012b;**25**:474–480.
- Dal Canto M, Guglielmo MC, Mignini Renzini M, Fadini R, Moutier C, Merola M, De Ponti E, Coticchio G. Dysmorphic patterns are associated with cytoskeletal alterations in human oocytes. *Hum Reprod* 2017;**32**:750–757.
- Daya S, Kohut J, Gunby J, Younglai E. Influence of blood clots in the cumulus complex on oocyte fertilization and cleavage. *Hum Reprod* 1990;**5**:744–746.
- de Cássia SFR, de Almeida Ferreira Braga DP, Semião-Francisco L, Madaschi C, Iaconelli A Jr, Borges E Jr. Metaphase II human oocyte morphology: contributing factors and effects on fertilization potential and embryo developmental ability in ICSI cycles. *Fertil Steril* 2010;**94**:1115–1117.
- De Croo I, De Sutter P, Tilleman K. A stepwise approach to move from a cleavage-stage to a blastocyst-stage transfer policy for all patients in the IVF clinic. *Hum Reprod Open* 2020;**2020**:hoaa034.
- De Geyter C, Wyns C, Calhaz-Jorge C, de Mouzon J, Ferraretti AP, Kupka M, Nyboe Andersen A, Nygren KG, Goossens V. 20 years of the European IVF-monitoring Consortium registry: what have we learned? A comparison with registries from two other regions. *Hum Reprod* 2020;**35**:2832–2849.
- De los Santos MJ, Apter S, Coticchio G, Debrock S, Lundin K, Plancha CE, Prados F, Rienzi L, Verheyen G, Woodward B et al; ESHRE Guideline Group on Good Practice in IVF Labs. Revised guidelines for good practice in IVF laboratories (2015). *Hum Reprod* 2016;**31**:685–686.
- De los Santos MJ, Arroyo G, Busquet A, Calderón G, Cuadros J, Hurtado de Mendoza MV, Moragas M, Herrer R, Ortiz A, Pons C et al; ASEBIR Interest Group in Embryology. A multicenter prospective study to assess the effect of early cleavage on embryo quality, implantation, and live-birth rate. *Fertil Steril* 2014;**101**:981–987.
- De Munck N, Bayram A, Elkhatib I, Abdala A, El-Damen A, Armanz A, Melado L, Lawrenz B, Fatemi HM. Marginal differences in preimplantation morphokinetics between conventional IVF and ICSI in patients with preimplantation genetic testing for aneuploidy (PGT-A): A sibling oocyte study. *PLoS One* 2022;**17**:e0267241.
- De Santis L, Cino I, Rabbellotti E, Calzi F, Persico P, Borini A, Coticchio G. Polar body morphology and spindle imaging as predictors of oocyte quality. *Reprod Biomed Online* 2005;**11**:36–42.
- De Sutter P, Dozortsev D, Qian C, Dhont M. Oocyte morphology does not correlate with fertilization rate and embryo quality after intracytoplasmic sperm injection. *Hum Reprod* 1996;**11**:595–597.
- De Vos A, Van de Velde H, Joris H, Van Steirteghem A. In-vitro matured metaphase-I oocytes have a lower fertilization rate but similar embryo quality as mature metaphase-II oocytes after intracytoplasmic sperm injection. *Hum Reprod* 1999;**14**:1859–1863.
- De Vos A, Van Landuyt L, Santos-Ribeiro S, Camus M, Van de Velde H, Tourmaye H, Verheyen G. Cumulative live birth rates after fresh and vitrified cleavage-stage versus blastocyst-stage embryo transfer in the first treatment cycle. *Hum Reprod* 2016;**31**:2442–2449.
- Desai N, Gill P. Blastomere cleavage plane orientation and the tetrahedral formation are associated with increased probability of a good-quality blastocyst for cryopreservation or transfer: a time-lapse study. *Fertil Steril* 2019;**111**:1159–1168.e1.
- Desai N, Goldberg JM, Austin C, Falcone T. Are cleavage anomalies, multinucleation, or specific cell cycle kinetics observed with time-lapse imaging predictive of embryo developmental capacity or ploidy? *Fertil Steril* 2018;**109**:665–674.
- Desai N, Ploskonka S, Goodman LR, Austin C, Goldberg J, Falcone T. Analysis of embryo morphokinetics, multinucleation and

- cleavage anomalies using continuous time-lapse monitoring in blastocyst transfer cycles. *Reprod Biol Endocrinol* 2014;**12**:54.
- Desch L, Bruno C, Luu M, Barberet J, Choux C, Lamotte M, Schmutz E, Sagot P, Fauque P. Embryo multinucleation at the two-cell stage is an independent predictor of intracytoplasmic sperm injection outcomes. *Fertil Steril* 2017;**107**:97–103.e104.
- Destouni A, Dimitriadou E, Masset H, Debrock S, Melotte C, Van Den Bogaert K, Zamani Esteki M, Ding J, Voet T, Denayer E et al Genome-wide haplotyping embryos developing from OPN and 1PN zygotes increases transferrable embryos in PGT-M. *Hum Reprod* 2018;**33**:2302–2311.
- Diakiw SM, Hall JMM, VerMilyea M, Lim AYY, Quangkananurug W, Chanchamroen S, Bankowski B, Stones R, Storr A, Miller A et al An artificial intelligence model correlated with morphological and genetic features of blastocyst quality improves ranking of viable embryos. *Reprod Biomed Online* 2022;**45**:1105–1117.
- Dietrich JE, Freis A, Beedgen F, von Horn K, Holschbach V, Liebscher J, Strowitzki T, Germeyer A. Intraindividual embryo morphokinetics are not affected by a switch of the ovarian stimulation protocol between GnRH agonist vs. antagonist regimens in consecutive cycles. *Front Endocrinol (Lausanne)* 2020;**11**:246.
- Dirican EK, Olgan S, Sakinci M, Caglar M. Blastocyst versus cleavage transfers: who benefits? *Arch Gynecol Obstet* 2022;**305**:749–756.
- Du QY, Wang EY, Huang Y, Guo XY, Xiong YJ, Yu YP, Yao GD, Shi SL, Sun YP. Blastocoele expansion degree predicts live birth after single blastocyst transfer for fresh and vitrified/warmed single blastocyst transfer cycles. *Fertil Steril* 2016;**105**:910–919.e1.
- Du T, Wang Y, Fan Y, Zhang S, Yan Z, Yu W, Xi Q, Chen Q, Mol BW, Lyu Q et al Fertility and neonatal outcomes of embryos achieving blastulation on Day 7: are they of clinical value? *Hum Reprod* 2018;**33**:1038–1051.
- Eastick J, Venetis C, Cooke S, Chapman M. The presence of cytoplasmic strings in human blastocysts is associated with the probability of clinical pregnancy with fetal heart. *J Assist Reprod Genet* 2021;**38**:2139–2149.
- Eastick J, Venetis C, Cooke S, Chapman M. Detailed analysis of cytoplasmic strings in human blastocysts: new insights. *Zygote* 2023a;**31**:78–84.
- Eastick J, Venetis C, Cooke S, Chapman M. Inter- and intra-observer agreement between embryologists for cytoplasmic string assessment in day 5/6 human blastocysts. *Reprod Sci* 2023b;**30**:1917–1926.
- Ebner T, Höggerl A, Oppelt P, Radler E, Enzelsberger SH, Mayer RB, Petek E, Shebl O. Time-lapse imaging provides further evidence that planar arrangement of blastomeres is highly abnormal. *Arch Gynecol Obstet* 2017;**296**:1199–1205.
- Ebner T, Maurer M, Shebl O, Moser M, Mayer RB, Duba HC, Tews G. Planar embryos have poor prognosis in terms of blastocyst formation and implantation. *Reprod Biomed Online* 2012;**25**:267–272.
- Ebner T, Moser M, Shebl O, Sommergruber M, Gaiswinkler U, Tews G. Morphological analysis at compacting stage is a valuable prognostic tool for ICSI patients. *Reprod Biomed Online* 2009;**18**:61–66.
- Ebner T, Moser M, Shebl O, Sommergruber M, Yaman C, Tews G. Blood clots in the cumulus-oocyte complex predict poor oocyte quality and post-fertilization development. *Reprod Biomed Online* 2008a;**16**:801–807.
- Ebner T, Moser M, Shebl O, Sommergruber M, Tews G. Prognosis of oocytes showing aggregation of smooth endoplasmic reticulum. *Reprod Biomed Online* 2008b;**16**:113–118.
- Ebner T, Moser M, Sommergruber M, Gaiswinkler U, Shebl O, Jesacher K, Tews G. Occurrence and developmental consequences of vacuoles throughout preimplantation development. *Fertil Steril* 2005;**83**:1635–1640.
- Ebner T, Moser M, Sommergruber M, Gaiswinkler U, Wiesinger R, Puchner M, Tews G. Presence, but not type or degree of extension, of a cytoplasmic halo has a significant influence on preimplantation development and implantation behaviour. *Hum Reprod* 2003;**18**:2406–2412.
- Ebner T, Moser M, Tews G. Is oocyte morphology prognostic of embryo developmental potential after ICSI? *Reprod Biomed Online* 2006;**12**:507–512.
- Ebner T, Sesli Ö, Kresic S, Enengl S, Stoiber B, Reiter E, Oppelt P, Mayer RB, Shebl O. Time-lapse imaging of cytoplasmic strings at the blastocyst stage suggests their association with spontaneous blastocoele collapse. *Reprod Biomed Online* 2020;**40**:191–199.
- Ebner T, Shebl O, Moser M, Sommergruber M, Tews G. Developmental fate of ovoid oocytes. *Hum Reprod* 2008c;**23**:62–66.
- Ebner T, Tritscher K, Mayer RB, Oppelt P, Duba HC, Maurer M, Schappacher-Tilp G, Petek E, Shebl O. Quantitative and qualitative trophectoderm grading allows for prediction of live birth and gender. *J Assist Reprod Genet* 2016;**33**:49–57.
- Ebner T, Yaman C, Moser M, Sommergruber M, Feichtinger O, Tews G. Prognostic value of first polar body morphology on fertilization rate and embryo quality in intracytoplasmic sperm injection. *Hum Reprod* 2000;**15**:427–430.
- Ebner T, Yaman C, Moser M, Sommergruber M, Pölz W, Tews G. Embryo fragmentation in vitro and its impact on treatment and pregnancy outcome. *Fertil Steril* 2001;**76**:281–285.
- Edwards RG, Steptoe PC, Purdy JM. Establishing full-term human pregnancies using cleaving embryos grown in vitro. *Br J Obstet Gynaecol* 1980;**87**:737–756.
- Ergin EG, Calışkan E, Yalçinkaya E, Oztel Z, Cökelez K, Ozay A, Özörnek HM. Frequency of embryo multinucleation detected by time-lapse system and its impact on pregnancy outcome. *Fertil Steril* 2014;**102**:1029–1033.e1.
- Escrich L, Galiana Y, Grau N, Insua F, Soler N, Pellicer A, Escrivá MJ. Do immature and mature sibling oocytes recovered from stimulated cycles have the same reproductive potential? *Reprod Biomed Online* 2018;**37**:667–676.
- Esfandiari N, Burjaq H, Gottlieb L, Casper RF. Brown oocytes: implications for assisted reproductive technology. *Fertil Steril* 2006;**86**:1522–1525.
- ESHRE Campus Course Report. Prevention of twin pregnancies after IVF/ICSI by single embryo transfer. *Hum Reprod* 2001;**16**:790–800.
- ESHRE Clinic PI Working Group, Vlaisavljevic V, Apter S, Capalbo A, D'Angelo A, Gianaroli L, Griesinger G, Kolibianakis EM, Lainas G, Mardesic T, Motrenko T. The Maribor consensus: report of an expert meeting on the development of performance indicators for clinical practice in ART. *Hum Reprod Open* 2021;**2021**:hoab022.
- ESHRE guideline group on the number of embryos to transfer. ESHRE Guideline: number of embryos to transfer during IVF/ICSI. *Hum Reprod* 2024;**39**:647–657.
- ESHRE PGT Consortium and SIG-Embryology Biopsy Working Group. ESHRE PGT Consortium and SIG Embryology good practice recommendations for polar body and embryo biopsy for PGT. *Human Reproduction Open* 2020;**2020**:hoaa020.
- ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine. The Vienna consensus: report of an expert meeting on the development of art laboratory performance indicators. *Hum Reprod Open* 2017;**2017**:hox011.
- ESHRE Working group on Time-lapse technology, Apter S, Ebner T, Freour T, Guns Y, Kovacic B, Le Clef N, Marques M, Meseguer M, Montjean D, Sfountouris I. Good practice recommendations for the use of time-lapse technology. *Hum Reprod Open* 2020;**2020**:hoaa008.
- Ezoe K, Coticchio G, Takenouchi H, Taoda S, Namerikawa S, Honda K, Miki T, Okimura T, Kobayashi T, Borini A et al Spatiotemporal

- perturbations of pronuclear breakdown preceding syngamy affect early human embryo development: a retrospective observational study. *J Assist Reprod Genet* 2022a;**39**:75–84.
- Ezoe K, Hickman C, Miki T, Okimura T, Uchiyama K, Yabuuchi A, Kobayashi T, Coticchio G, Kato K. Cytoplasmic halo characteristics during fertilization and their implications for human preimplantation embryo development and pregnancy outcome. *Reprod Biomed Online* 2020;**41**:191–202.
- Ezoe K, Miki T, Akaike H, Shimazaki K, Takahashi T, Tanimura Y, Amagai A, Sawado A, Mogi M, Kaneko S et al Maternal age affects pronuclear and chromatin dynamics, morula compaction and cell polarity, and blastulation of human embryos. *Hum Reprod* 2023;**38**:387–399.
- Ezoe K, Shimazaki K, Miki T, Takahashi T, Tanimura Y, Amagai A, Sawado A, Akaike H, Mogi M, Kaneko S et al Association between a deep learning-based scoring system with morphokinetics and morphological alterations in human embryos. *Reprod Biomed Online* 2022b;**45**:1124–1132.
- Ezoe K, Takahashi T, Shimazaki K, Miki T, Tanimura Y, Amagai A, Sawado A, Akaike H, Mogi M, Kaneko S et al Human 1PN and 3PN zygotes recapitulate all morphokinetic events of normal fertilization but reveal novel developmental errors. *Hum Reprod* 2022c;**37**:2307–2319.
- Fabozzi G, Alteri A, Rega E, Starita MF, Piscitelli C, Giannini P, Colicchia A. Morphological assessment on day 4 and its prognostic power in selecting viable embryos for transfer. *Zygote* 2016;**24**:477–484.
- Fancsovits P, Murber A, Gilán ZT, Rigó J Jr, Urbancsek J. Human oocytes containing large cytoplasmic vacuoles can result in pregnancy and viable offspring. *Reprod Biomed Online* 2011;**23**:513–516.
- Fancsovits P, Tóthné ZG, Murber A, Takács FZ, Papp Z, Urbancsek J. Correlation between first polar body morphology and further embryo development. *Acta Biol Hung* 2006;**57**:331–338.
- Fang T, Yu W, Ou S, Lu J, Li R, Zhao M, Chan YL, Wang W. The impact of oocytes containing smooth endoplasmic reticulum aggregates on assisted reproductive outcomes: a cohort study. *BMC Pregnancy Childbirth* 2022;**22**:838.
- Faramarzi A, Khalili MA, Ashourzadeh S. Oocyte morphology and embryo morphokinetics in an intra-cytoplasmic sperm injection programme. Is there a relationship? *Zygote* 2017;**25**:190–196.
- Faramarzi A, Khalili MA, Ashourzadeh S, Palmerini MG. Does rescue in vitro maturation of germinal vesicle stage oocytes impair embryo morphokinetics development? *Zygote* 2018;**26**:430–434.
- Faramarzi A, Khalili MA, Omid M. Morphometric analysis of human oocytes using time lapse: does it predict embryo developmental outcomes? *Hum Fertil (Camb)* 2019;**22**:171–176.
- Farhi J, Nahum H, Weissman A, Zahalka N, Glezerman M, Levran D. Coarse granulation in the perivitelline space and IVF-ICSI outcome. *J Assist Reprod Genet* 2002;**19**:545–549.
- Feil D, Henshaw RC, Lane M. Day 4 embryo selection is equal to Day 5 using a new embryo scoring system validated in single embryo transfers. *Hum Reprod* 2008;**23**:1505–1510.
- Ferrari Zanetti B, Paes de Almeida Ferreira Braga D, Souza Setti A, de Cássia Sávio Figueira R, Iaconelli A Jr, Borges E Jr. Is perivitelline space morphology of the oocyte associated with pregnancy outcome in intracytoplasmic sperm injection cycles? *Eur J Obstet Gynecol Reprod Biol* 2018;**231**:225–229.
- Firmin J, Ecker N, Rivet Danon D, Özgüç Ö, Barraud Lange V, Turlier H, Patrat C, Maitre JL. Mechanics of human embryo compaction. *Nature* 2024;**629**:646–651.
- Fishel S, Campbell A, Foad F, Davies L, Best L, Davis N, Smith R, Duffy S, Wheat S, Montgomery S et al Evolution of embryo selection for IVF from subjective morphology assessment to objective time-lapse algorithms improves chance of live birth. *Reprod Biomed Online* 2020;**40**:61–70.
- Fishel S, Campbell A, Montgomery S, Smith R, Nice L, Duffy S, Jenner L, Berrisford K, Kellam L, Smith R et al Time-lapse imaging algorithms rank human preimplantation embryos according to the probability of live birth. *Reprod Biomed Online* 2018;**37**:304–313.
- Fitz VW, Kanakasabapathy MK, Thirumalaraju P, Kandula H, Ramirez LB, Boehnlein L, Swain JE, Curchoe CL, James K, Dimitriadis I et al Should there be an “AI” in TEAM? Embryologists selection of high implantation potential embryos improves with the aid of an artificial intelligence algorithm. *J Assist Reprod Genet* 2021;**38**:2663–2670.
- Forman EJ, Upham KM, Cheng M, Zhao T, Hong KH, Treff NR, Scott RT Jr. Comprehensive chromosome screening alters traditional morphology-based embryo selection: a prospective study of 100 consecutive cycles of planned fresh euploid blastocyst transfer. *Fertil Steril* 2013;**100**:718–724.
- Fouks Y, Yogev Y. Twinning in ART: single embryo transfer policy. *Best Pract Res Clin Obstet Gynaecol* 2022;**84**:88–95.
- Fraire-Zamora JJ, Serdarogullari M, Sharma K, Ammar OF, Mincheva M, Macklon N, Pujol A, Capalbo A, Meseguer M, Liperis G. Better late than never: the clinical value of Day 7 blastocysts. *Hum Reprod* 2023;**38**:520–524.
- Fu L, Chen S, Wang M, Huang G, Wang F, Lan Y, Liu S, Jiang X. Live birth from a blastocyst derived from a conjoined oocyte in a frozen embryo transfer cycle: a case report and a literature review. *J Assist Reprod Genet* 2022a;**39**:1351–1357.
- Fu L, Chu D, Zhou W, Li Y. Strictly selected mono- and non-pronuclear blastocysts could result in appreciable clinical outcomes in IVF cycles. *Hum Fertil (Camb)* 2022b;**25**:470–477.
- Fu L, Zhou W, Li Y. Development and frozen-thawed transfer of non-pronuclear zygotes-derived embryos in IVF cycles. *Eur J Obstet Gynecol Reprod Biol* 2021;**264**:206–211.
- Gardner DK. The impact of physiological oxygen during culture, and vitrification for cryopreservation, on the outcome of extended culture in human IVF. *Reprod Biomed Online* 2016;**32**:137–141.
- Gardner DK, Kelley RL. Impact of the IVF laboratory environment on human preimplantation embryo phenotype. *J Dev Orig Health Dis* 2017;**8**:418–435.
- Gardner DK, Lane M. Alleviation of the ‘2-cell block’ and development to the blastocyst of CF1 mouse embryos: role of amino acids, EDTA and physical parameters. *Hum Reprod* 1996;**11**:2703–2712.
- Gardner DK, Lane M. Culture and selection of viable blastocysts: a feasible proposition for human IVF? *Hum Reprod Update* 1997;**3**:367–382.
- Gardner DK, Lane M. Culture Systems for the Human Embryo. In: Gardner DK, Weissman A, Howles C, Shoham Z (eds). *Textbook of Assisted Reproductive Techniques*, 5th edn. Boca Raton, USA: CRC Press, 2018, 200–224.
- Gardner DK, Schoolcraft WB. In-vitro culture of human blastocysts. In: Jansen R, Mortimer D (eds). *Towards Reproductive Certainty: Fertility and Genetics Beyond 1999*. Carnforth, UK: Parthenon Press, 1999, 378–388.
- Ghosh J, Coutifaris C, Sapienza C, Mainigi M. Global DNA methylation levels are altered by modifiable clinical manipulations in assisted reproductive technologies. *Clin Epigenetics* 2017;**9**:14.
- Giorgetti C, Terriou P, Auquier P, Hans E, Spach JL, Salzmann J, Roulier R. Embryo score to predict implantation after in-vitro fertilization: based on 957 single embryo transfers. *Hum Reprod* 1995;**10**:2427–2431.
- Goodman LR, Goldberg J, Falcone T, Austin C, Desai N. Does the addition of time-lapse morphokinetics in the selection of embryos for

- transfer improve pregnancy rates? A randomized controlled trial. *Fertil Steril* 2016;**105**:275–285.e10.
- Grøndahl ML, Christiansen SL, Kesmodel US, Agerholm IE, Lemmen JG, Lundstrøm P, Bogstad J, Raaschou-Jensen M, Ladelund S. Effect of women's age on embryo morphology, cleavage rate and competence—a multicenter cohort study. *PLoS One* 2017;**12**:e0172456.
- Gurunath S, Biliangady R, Sundhararaj UM, Gangadharswamy A, Gundlapalli S, Reddy GMM. Live birth rates in in vitro fertilization cycles with oocytes containing smooth endoplasmic reticulum aggregates and normal oocytes. *J Hum Reprod Sci* 2019;**12**:156–163.
- Halim B, Lubis HP, Adenin I, Angellee J, Samoedra RS. Meiotic spindle view improves the outcome of IVF in poor responders: a retrospective analytical study from an Indonesian IVF Center. *JBRA Assist Reprod* 2024;**28**:39–46.
- Hammond ER, Cree LM, Morbeck DE. Should extended blastocyst culture include Day 7? *Hum Reprod* 2018;**33**:991–997.
- Hammond ER, Foong AKM, Rosli N, Morbeck DE. Should we freeze it? Agreement on fate of borderline blastocysts is poor and does not improve with a modified blastocyst grading system. *Hum Reprod* 2020;**35**:1045–1053.
- Harbottle S, Hughes C, Cutting R, Roberts S, Brison D; Association Of Clinical Embryologists (ACE) & The British Fertility Society (BFS). Elective single embryo transfer: an update to UK Best Practice Guidelines. *Hum Fertil (Camb)* 2015;**18**:165–183.
- Hardarson T, Hanson C, Sjögren A, Lundin K. Human embryos with unevenly sized blastomeres have lower pregnancy and implantation rates: indications for aneuploidy and multinucleation. *Hum Reprod* 2001;**16**:313–318.
- Hassan-Ali H, Hisham-Saleh A, El-Gezeiry D, Baghdady I, Ismaeil I, Mandelbaum J. Perivitelline space granularity: a sign of human menopausal gonadotrophin overdose in intracytoplasmic sperm injection. *Hum Reprod* 1998;**13**:3425–3430.
- Hattori H, Nakamura Y, Nakajo Y, Araki Y, Kyono K. Deliveries of babies with normal health derived from oocytes with smooth endoplasmic reticulum clusters. *J Assist Reprod Genet* 2014;**31**:1461–1467.
- Herrero J, Tejera A, Albert C, Vidal C, de los Santos MJ, Meseguer M. A time to look back: analysis of morphokinetic characteristics of human embryo development. *Fertil Steril* 2013;**100**:1602–1609.
- Hill MJ, Richter KS, Heitmann RJ, Graham JR, Tucker MJ, DeCherney AH, Browne PE, Levens ED. Trophoctoderm grade predicts outcomes of single-blastocyst transfers. *Fertil Steril* 2013;**99**:1283–1289.e1281.
- Hoek J, Schoenmakers S, van Duijn L, Willemsen SP, van Marion ES, Laven JSE, Baart EB, Steegers-Theunissen RPM. A higher pre-conceptional paternal body mass index influences fertilization rate and preimplantation embryo development. *Andrology* 2022;**10**:486–494.
- Holte J, Berglund L, Milton K, Garello C, Gennarelli G, Revelli A, Bergh T. Construction of an evidence-based integrated morphology cleavage embryo score for implantation potential of embryos scored and transferred on day 2 after oocyte retrieval. *Hum Reprod* 2007;**22**:548–557.
- Holubcová Z, Kyjovská D, Martonová M, Páralová D, Klenková T, Otevřel P, Štěpánová R, Kloudová S, Hampl A. Egg maturity assessment prior to ICSI prevents premature fertilization of late-maturing oocytes. *J Assist Reprod Genet* 2019;**36**:445–452.
- Hondo S, Arichi A, Muramatsu H, Omura N, Ito K, Komine H, Monzen S, Mukai N, Endo M, Katase S et al. Clinical outcomes of transfer of frozen and thawed single blastocysts derived from nonpronuclear and monopronuclear zygotes. *Reprod Med Biol* 2019;**18**:278–283.
- Honnma H, Baba T, Sasaki M, Hashiba Y, Ohno H, Fukunaga T, Endo T, Saito T, Asada Y. Trophoctoderm morphology significantly affects the rates of ongoing pregnancy and miscarriage in frozen-thawed single-blastocyst transfer cycle in vitro fertilization. *Fertil Steril* 2012;**98**:361–367.
- Hori K, Hori K, Kosasa T, Walker B, Ohta A, Ahn HJ, Huang TTF. Comparison of euploid blastocyst expansion with subgroups of single chromosome, multiple chromosome, and segmental aneuploids using an AI platform from donor egg embryos. *J Assist Reprod Genet* 2023;**40**:1407–1416.
- Huang TT, Huang DH, Ahn HJ, Arnett C, Huang CT. Early blastocyst expansion in euploid and aneuploid human embryos: evidence for a non-invasive and quantitative marker for embryo selection. *Reprod Biomed Online* 2019;**39**:27–39.
- Huang TTF, Kosasa T, Walker B, Arnett C, Huang CTF, Yin C, Harun Y, Ahn HJ, Ohta A. Deep learning neural network analysis of human blastocyst expansion from time-lapse image files. *Reprod Biomed Online* 2021;**42**:1075–1085.
- Hung TY, Lee RK, Hwu YM, Lin MH, Li RS, Weng YW. Early blastulation of day 4 embryo correlates with the increased euploid rate of preimplantation genetic screening cycles. *Taiwan J Obstet Gynecol* 2018;**57**:858–861.
- Hur C, Nanavaty V, Yao M, Desai N. The presence of partial compaction patterns is associated with lower rates of blastocyst formation, sub-optimal morphokinetic parameters and poorer morphologic grade. *Reprod Biol Endocrinol* 2023;**21**:12.
- Hviid KVR, Malchau SS, Pinborg A, Nielsen HS. Determinants of monozygotic twinning in ART: a systematic review and a meta-analysis. *Hum Reprod Update* 2018;**24**:468–483.
- Illingworth PJ, Venetis C, Gardner DK, Nelson SM, Berntsen J, Larman MG, Agresta F, Ahitan S, Ahlström A, Cattrall F et al. Deep learning versus manual morphology-based embryo selection in IVF: a randomized, double-blind noninferiority trial. *Nat Med* 2024;**30**:3114–3120.
- Inoue T, Taguchi S, Uemura M, Tsujimoto Y, Kokunai K, Ikawa K, Yamashita Y. The migration speed of nucleolar precursor bodies in pronuclei affects in vitro fertilization-derived human embryo ploidy status and live birth. *Reprod Med Biol* 2023;**22**:e12497.
- Inoue T, Taguchi S, Uemura M, Tsujimoto Y, Miyazaki K, Yamashita Y. Migration speed of nucleolus precursor bodies in human male pronuclei: a novel parameter for predicting live birth. *J Assist Reprod Genet* 2021;**38**:1725–1736.
- Irani M, Reichman D, Robles A, Melnick A, Davis O, Zaninovic N, Xu K, Rosenwaks Z. Morphologic grading of euploid blastocysts influences implantation and ongoing pregnancy rates. *Fertil Steril* 2017;**107**:664–670.
- Itoi F, Asano Y, Shimizu M, Honnma H, Murata Y. Birth of nine normal healthy babies following transfer of blastocysts derived from human single-pronucleate zygotes. *J Assist Reprod Genet* 2015;**32**:1401–1407.
- Itoi F, Asano Y, Shimizu M, Nagai R, Saitou K, Honnma H, Murata Y. Clinical outcomes after IVF or ICSI using human blastocysts derived from oocytes containing aggregates of smooth endoplasmic reticulum. *Reprod Biomed Online* 2017;**34**:337–344.
- Ivec M, Kovacic B, Vlaisavljevic V. Prediction of human blastocyst development from morulas with delayed and/or incomplete compaction. *Fertil Steril* 2011;**96**:1473–1478.e2.
- Iwata K, Yumoto K, Sugishima M, Mizoguchi C, Kai Y, Iba Y, Mio Y. Analysis of compaction initiation in human embryos by using time-lapse cinematography. *J Assist Reprod Genet* 2014;**31**:421–426.
- Jackson KV, Ginsburg ES, Hornstein MD, Rein MS, Clarke RN. Multinucleation in normally fertilized embryos is associated with an accelerated ovulation induction response and lower

- implantation and pregnancy rates in in vitro fertilization-embryo transfer cycles. *Fertil Steril* 1998;**70**:60–66.
- Ji M, Wang X, Wu W, Guan Y, Liu J, Wang J, Liu W, Shen C. ART manipulation after controlled ovarian stimulation may not increase the risk of abnormal expression and DNA methylation at some CpG sites of H19, IGF2 and SNRPN in fetuses: a pilot study. *Reprod Biol Endocrinol* 2018;**16**:63.
- Johansson M, Hardarson T, Lundin K. There is a cutoff limit in diameter between a blastomere and a small anucleate fragment. *J Assist Reprod Genet* 2003;**20**:309–313.
- Joo K, Nemes A, Dudas B, Berkes-Bara E, Murber A, Urbancsek J, Fancsovits P. The importance of cytoplasmic strings during early human embryonic development. *Front Cell Dev Biol* 2023;**11**:1177279.
- Kadam N, Woodhead G, Kellam L, Campbell A, Jayaprakasan K. Odds and predictors of monozygotic twinning in a multicentre cohort of 25,794 IVF cycles. *JCM* 2023;**12**:2593.
- Kahraman S, Yakin K, Dönmez E, Samli H, Bahçe M, Cengiz G, Sertyel S, Samli M, Imirzalioglu N. Relationship between granular cytoplasm of oocytes and pregnancy outcome following intracytoplasmic sperm injection. *Hum Reprod* 2000;**15**:2390–2393.
- Kai Y, Iwata K, Iba Y, Mio Y. Diagnosis of abnormal human fertilization status based on pronuclear origin and/or centrosome number. *J Assist Reprod Genet* 2015;**32**:1589–1595.
- Kai Y, Moriwaki H, Yumoto K, Iwata K, Mio Y. Assessment of developmental potential of human single pronucleated zygotes derived from conventional in vitro fertilization. *J Assist Reprod Genet* 2018;**35**:1377–1384.
- Kato K, Ueno S, Berntsen J, Kragh MF, Okimura T, Kuroda T. Does embryo categorization by existing artificial intelligence, morphokinetic or morphological embryo selection models correlate with blastocyst euploidy rates? *Reprod Biomed Online* 2023;**46**:274–281.
- Kemper JM, Liu Y, Afnan M, Hammond ER, Morbeck DE, Mol BW. Should we look for a low-grade threshold for blastocyst transfer? A scoping review. *Reprod Biomed Online* 2021;**42**:709–716.
- Kemper JM, Liu Y, Afnan M, Mol BWJ, Morbeck DE. What happens to abnormally fertilized embryos? A scoping review. *Reprod Biomed Online* 2023;**46**:802–807.
- Khosravi P, Kazemi E, Zhan Q, Malmsten JE, Toschi M, Zisimopoulos P, Sigaras A, Lavery S, Cooper LAD, Hickman C et al Deep learning enables robust assessment and selection of human blastocysts after in vitro fertilization. *NPJ Digit Med* 2019;**2**:21.
- Kieslinger DC, Vergouw CG, Ramos L, Arends B, Curfs M, Slappendel E, Kosteljik EH, Pieters M, Consten D, Verhoeven MO et al Clinical outcomes of uninterrupted embryo culture with or without time-lapse-based embryo selection versus interrupted standard culture (SelecTIMO): a three-armed, multicentre, double-blind, randomised controlled trial. *Lancet* 2023;**401**:1438–1446.
- Kirkegaard K, Hindkjaer JJ, Ingerslev HJ. Effect of oxygen concentration on human embryo development evaluated by time-lapse monitoring. *Fertil Steril* 2013;**99**:738–744.e734.
- Kirkegaard K, Sundvall L, Erlandsen M, Hindkjær JJ, Knudsen UB, Ingerslev HJ. Timing of human preimplantation embryonic development is confounded by embryo origin. *Hum Reprod* 2016;**31**:324–331.
- Kitasaka H, Konuma Y, Tokoro M, Fukunaga N, Asada Y. Oocyte cytoplasmic diameter of $\geq 130 \mu\text{m}$ can be used to determine human giant oocytes. *F S Sci* 2022;**3**:10–17.
- Kligman I, Benadiva C, Alikani M, Munne S. The presence of multinucleated blastomeres in human embryos is correlated with chromosomal abnormalities. *Hum Reprod* 1996;**11**:1492–1498.
- Kljajic M, Saymé N, Krebs T, Wagenpfeil G, Baus S, Solomayer EF, Kasoha M. Zygote diameter and total cytoplasmic volume as useful predictive tools of blastocyst quality. *Geburtshilfe Frauenheilkd* 2023;**83**:97–105.
- Knez J, Kovačić B, Vlasisavljević V. Comparison of embryo transfer strategies and assisted reproduction outcome in Slovenian and cross-border patients. *Reprod Biomed Online* 2013;**27**:310–315.
- Kong X, Yang S, Gong F, Lu C, Zhang S, Lu G, Lin G. The relationship between cell number, division behavior and developmental potential of cleavage stage human embryos: a time-lapse study. *PLoS One* 2016;**11**:e0153697.
- Kovačić B. Culture systems: low-oxygen culture. *Methods Mol Biol* 2012;**912**:249–272.
- Kovačić B. Incubators for embryo culture. In: Ahlström A, Lundin K (eds). *Manual of Embryo Culture in Human Assisted Reproduction*. Cambridge: Cambridge University Press, 2021, 7–19.
- Kovacic B, Vlasisavljevic V, Reljic M, Cizek-Sajko M. Developmental capacity of different morphological types of day 5 human morulae and blastocysts. *Reprod Biomed Online* 2004;**8**:687–694.
- Kramer YG, Kofinas JD, Melzer K, Noyes N, McCaffrey C, Buldo-Licciardi J, McCulloh DH, Grifo JA. Assessing morphokinetic parameters via time lapse microscopy (TLM) to predict euploidy: are aneuploidy risk classification models universal? *J Assist Reprod Genet* 2014;**31**:1231–1242.
- Kroener LL, Ambartsumyan G, Pisarska MD, Briton-Jones C, Surrey M, Hill D. Increased blastomere number in cleavage-stage embryos is associated with higher aneuploidy. *Fertil Steril* 2015;**103**:694–698.
- La Sala GB, Nicoli A, Villani MT, Di Girolamo R, Capodanno F, Blickstein I. The effect of selecting oocytes for insemination and transferring all resultant embryos without selection on outcomes of assisted reproduction. *Fertil Steril* 2009;**91**:96–100.
- Lagalla C, Coticchio G, Sciajno R, Tarozzi N, Zacà C, Borini A. Alternative patterns of partial embryo compaction: prevalence, morphokinetic history and possible implications. *Reprod Biomed Online* 2020;**40**:347–354.
- Lagalla C, Tarozzi N, Sciajno R, Wells D, Di Santo M, Nadalini M, Distratis V, Borini A. Embryos with morphokinetic abnormalities may develop into euploid blastocysts. *Reprod Biomed Online* 2017;**34**:137–146.
- Lane SL, Reed L, Schoolcraft WB, Katz-Jaffe MG. Euploid day 7 blastocysts of infertility patients with only slow embryo development have reduced implantation potential. *Reprod Biomed Online* 2022;**44**:858–865.
- Langley MT, Marek DM, Gardner DK, Doody KM, Doody KJ. Extended embryo culture in human assisted reproduction treatments. *Hum Reprod* 2001;**16**:902–908.
- Le Cruguel S, Ferré-L'Hôtelier V, Morinière C, Lemerle S, Reynier P, Descamps P, May-Panloup P. Early compaction at day 3 may be a useful additional criterion for embryo transfer. *J Assist Reprod Genet* 2013;**30**:683–690.
- Lebovitz O, Michaeli M, Aslih N, Poltov D, Estrada D, Atzmon Y, Shalom-Paz E. Embryonic development in relation to maternal age and conception probability. *Reprod Sci* 2021;**28**:2292–2300.
- Lehner A, Kaszas Z, Murber A, Rigo J Jr, Urbancsek J, Fancsovits P. Giant oocytes in human in vitro fertilization treatments. *Arch Gynecol Obstet* 2015;**292**:697–703.
- Li M, Dang Y, Wang Y, Li J, Liu P. Value of transferring embryos derived from monopronucleated (1PN) zygotes at the time of fertilization assessment. *Zygote* 2020;**28**:241–246.
- Li M, Huang J, Zhuang X, Lin S, Dang Y, Wang Y, Liu D, Li R, Liu P, Qiao J. Obstetric and neonatal outcomes after the transfer of vitrified-warmed blastocysts developing from nonpronuclear and monopronuclear zygotes: a retrospective cohort study. *Fertil Steril* 2021;**115**:110–117.

- Licciardi F, McCaffrey C, Oh C, Schmidt-Sarosi C, McCulloh DH. Birth weight is associated with inner cell mass grade of blastocysts. *Fertil Steril* 2015;**103**:382–387. e382.
- Lin YC, Chang SY, Lan KC, Huang HW, Chang CY, Tsai MY, Kung FT, Huang FJ. Human oocyte maturity in vivo determines the outcome of blastocyst development in vitro. *J Assist Reprod Genet* 2003;**20**:506–512.
- Liu J, Wang XL, Zhang X, Shen CY, Zhang Z. Live births resulting from OPN-derived embryos in conventional IVF cycles. *J Assist Reprod Genet* 2016;**33**:373–378.
- Liu Y, Chapple V, Roberts P, Ali J, Matson P. Time-lapse videography of human oocytes following intracytoplasmic sperm injection: events up to the first cleavage division. *Reprod Biol* 2014;**14**:249–256.
- Liu Y, Peng X, Liu C, Zhang S, Weng Z, Yu L, Zhou S, Huang X. Live birth derived from a markedly large polar body oocyte: a rare case report. *Zygote* 2024;**32**:170–174.
- Liu Z, Jiang M, He L, Liu Y. Cell number considerations for blastocyst transfer in younger patients. *J Assist Reprod Genet* 2020;**37**:619–627.
- Loutradis D, Drakakis P, Kallianidis K, Milingos S, Dendrinis S, Michalas S. Oocyte morphology correlates with embryo quality and pregnancy rate after intracytoplasmic sperm injection. *Fertil Steril* 1999;**72**:240–244.
- Lundin K, Ahlström A. Quality control and standardization of embryo morphology scoring and viability markers. *Reprod Biomed Online* 2015;**31**:459–471.
- Lundin K, Bergh C, Hardarson T. Early embryo cleavage is a strong indicator of embryo quality in human IVF. *Hum Reprod* 2001;**16**:2652–2657.
- Ma BX, Yang L, Tian Y, Jin L, Huang B. Cytoplasmic strings between ICM and mTE are a positive predictor of clinical pregnancy and live birth outcomes: a time-lapse study. *Front Med (Lausanne)* 2022;**9**:934327.
- Machtinger R, Politch JA, Hornstein MD, Ginsburg ES, Racowsky C. A giant oocyte in a cohort of retrieved oocytes: does it have any effect on the in vitro fertilization cycle outcome? *Fertil Steril* 2011;**95**:573–576.
- Machtinger R, Racowsky C. Morphological systems of human embryo assessment and clinical evidence. *Reprod Biomed Online* 2013;**26**:210–221.
- Madaschi C, Aoki T, de Almeida Ferreira Braga DP, de Cássia Sávio Figueira R, Semião Francisco L, Iaconelli A Jr, Borges E Jr. Zona pellucida birefringence score and meiotic spindle visualization in relation to embryo development and ICSI outcomes. *Reprod Biomed Online* 2009;**18**:681–686.
- Magdi Y. Dizygotic twin from conjoined oocytes: a case report. *J Assist Reprod Genet* 2020;**37**:1367–1370.
- Magli MC, Gianaroli L, Ferraretti AP, Lappi M, Ruberti A, Farfalli V. Embryo morphology and development are dependent on the chromosomal complement. *Fertil Steril* 2007;**87**:534–541.
- Magli M C, Jones G M, Lundin K, Van Den Abbeel E. The Atlas of Human Embryology: from Oocytes to Preimplantation Embryos. Preface. *Hum Reprod* 2012;**27**:i1–i11.
- Marconi N, Raja EA, Bhattacharya S, Maheshwari A. Perinatal outcomes in singleton live births after blastocyst transfer: an analysis of 60,926 in vitro fertilization cycles from the United Kingdom. *Fertil Steril* 2023;**120**:312–320.
- Marcos J, Pérez-Albalá S, Mifsud A, Molla M, Landeras J, Meseguer M. Collapse of blastocysts is strongly related to lower implantation success: a time-lapse study. *Hum Reprod* 2015;**30**:2501–2508.
- Margalit T, Ben-Haroush A, Garor R, Kotler N, Shefer D, Krasilnikov N, Tzabari M, Oron G, Shufaro Y, Sapir O. Morphokinetic characteristics of embryos derived from in-vitro-matured oocytes and their in-vivo-matured siblings after ovarian stimulation. *Reprod Biomed Online* 2019;**38**:7–11.
- Martínez-Granados L, Serrano M, González-Utor A, Ortíz N, Badajoz V, Olaya E, Prados N, Boada M, Castilla JA; Special Interest Group in Quality of ASEBIR (Spanish Society for the Study of Reproductive Biology). Inter-laboratory agreement on embryo classification and clinical decision: Conventional morphological assessment vs. time lapse. *PLoS One* 2017;**12**:e0183328.
- Martins WP, Nastro CO, Rienzi L, van der Poel SZ, Gracia CR, Racowsky C. Obstetrical and perinatal outcomes following blastocyst transfer compared to cleavage transfer: a systematic review and meta-analysis. *Hum Reprod* 2016;**31**:2561–2569.
- Massarotti C, Stigliani S, Ramone A, Bovis F, Sozzi F, Remorgida V, Cagnacci A, Anserini P, Scaruffi P. Occurrence of smooth endoplasmic reticulum aggregates in metaphase II oocytes: relationship with stimulation protocols and outcome of ICSI and IVF cycles. *Hum Reprod* 2021;**36**:907–917.
- Mateizel I, Van Landuyt L, Tournaye H, Verheyen G. Deliveries of normal healthy babies from embryos originating from oocytes showing the presence of smooth endoplasmic reticulum aggregates. *Hum Reprod* 2013;**28**:2111–2117.
- Mayer RB, Shebl O, Oppelt P, Reiter E, Altmann R, Enengl S, Allerstorfer C, Ebner T. Good-quality blastocysts derived from vacuolized morulas show reduced viability. *Fertil Steril* 2018;**109**:1025–1029.
- Meseguer M, Herrero J, Tejera A, Hilligsøe KM, Ramsing NB, Remohí J. The use of morphokinetics as a predictor of embryo implantation. *Hum Reprod* 2011;**26**:2658–2671.
- Meseguer M, Rubio I, Cruz M, Basile N, Marcos J, Requena A. Embryo incubation and selection in a time-lapse monitoring system improves pregnancy outcome compared with a standard incubator: a retrospective cohort study. *Fertil Steril* 2012;**98**:1481–1489. e10.
- Milewski R, Kuć P, Kuczyńska A, Stankiewicz B, Łukaszuk K, Kuczyński W. A predictive model for blastocyst formation based on morphokinetic parameters in time-lapse monitoring of embryo development. *J Assist Reprod Genet* 2015;**32**:571–579.
- Minasi MG, Colasante A, Riccio T, Ruberti A, Casciani V, Scarselli F, Spinella F, Fiorentino F, Varricchio MT, Greco E. Correlation between aneuploidy, standard morphology evaluation and morphokinetic development in 1730 biopsied blastocysts: a consecutive case series study. *Hum Reprod* 2016;**31**:2245–2254.
- Mio Y, Maeda K. Time-lapse cinematography of dynamic changes occurring during in vitro development of human embryos. *Am J Obstet Gynecol* 2008;**199**:660.e1–660.e5.
- Mizobe Y, Kuwatsuru Y, Kuroki Y, Fukumoto Y, Tokudome M, Moewaki H, Tabira M, Iwakawa T, Takeuchi K. Smooth endoplasmic reticulum cluster presence does not affect embryo ploidy. *Arch Gynecol Obstet* 2023;**307**:1607–1612.
- Montag M, Liebenthron J, Köster M. Which morphological scoring system is relevant in human embryo development? *Placenta* 2011;**32** Suppl 3:S252–S256.
- Montag M, Schimming T, Köster M, Zhou C, Dorn C, Rösing B, van der Ven H, Ven der Ven K. Oocyte zona birefringence intensity is associated with embryonic implantation potential in ICSI cycles. *Reprod Biomed Online* 2008;**16**:239–244.
- Montjean D, Geoffroy-Siraudin C, Gervoise-Boyer M-J, Boyer P. Competence of embryos showing transient developmental arrest during in vitro culture. *J Assist Reprod Genet* 2021;**38**:857–863.
- Moon JH, Zhao Q, Zhang J, Reddy V, Han J, Cheng Y, Zhang N, Dasig J, Nel-Themaat L, Behr B et al The developmental competence of human metaphase I oocytes with delayed maturation in vitro. *Fertil Steril* 2023;**119**:690–696.

- Morbeck DE. Blastocyst culture in the Era of PGS and FreezeAlls: Is a 'C' a failing grade? *Hum Reprod Open* 2017;**2017**:hox017.
- Motato Y, de los Santos MJ, Escriba MJ, Ruiz BA, Remohí J, Meseguer M. Morphokinetic analysis and embryonic prediction for blastocyst formation through an integrated time-lapse system. *Fertil Steril* 2016;**105**:376–384.e9.
- Múgica A, Martín M, Soydan E, Esbert M, Ballesteros A, Calderón G. Relationship between day-3 embryo blastomere symmetry and implantation rate. *Fertil Steril* 2008;**90**:S342–S343.
- Mumusoglu S, Yarali I, Bozdog G, Ozdemir P, Polat M, Sokmensuer LK, Yarali H. Time-lapse morphokinetic assessment has low to moderate ability to predict euploidy when patient- and ovarian stimulation-related factors are taken into account with the use of clustered data analysis. *Fertil Steril* 2017;**107**:413–421.e4.
- Munné S, Alikani M, Tomkin G, Grifo J, Cohen J. Embryo morphology, developmental rates, and maternal age are correlated with chromosome abnormalities. *Fertil Steril* 1995;**64**:382–391.
- Munné S, Sandalinas M, Magli C, Gianaroli L, Cohen J, Warburton D. Increased rate of aneuploid embryos in young women with previous aneuploid conceptions. *Prenat Diagn* 2004;**24**:638–643.
- Mutia K, Wiweko B, Iffanolida PA, Febri RR, Muna N, Riayati O, Jasirwan SO, Yuningsih T, Mansyur E, Hestiantoro A. The Frequency of Chromosomal Euploidy Among 3PN Embryos. *J Reprod Infertil* 2019;**20**:127–131.
- Nagy ZP, Janssenswillen C, Janssens R, De Vos A, Staessen C, Van de Velde H, Van Steirteghem AC. Timing of oocyte activation, pronucleus formation and cleavage in humans after intracytoplasmic sperm injection (ICSI) with testicular spermatozoa and after ICSI or in-vitro fertilization on sibling oocytes with ejaculated spermatozoa. *Hum Reprod* 1998;**13**:1606–1612.
- Navarro PA, de Araújo MM, de Araújo CM, Rocha M, dos Reis R, Martins W. Relationship between first polar body morphology before intracytoplasmic sperm injection and fertilization rate, cleavage rate, and embryo quality. *Int J Gynaecol Obstet* 2009;**104**:226–229.
- Nazem TG, Sekhon L, Lee JA, Overbey J, Pan S, Duke M, Briton-Jones C, Whitehouse M, Copperman AB, Stein DE. The correlation between morphology and implantation of euploid human blastocysts. *Reprod Biomed Online* 2019;**38**:169–176.
- Ng ST, Chang TH, Wu TC. Prediction of the rates of fertilization, cleavage, and pregnancy success by cumulus-coronal morphology in an in vitro fertilization program. *Fertil Steril* 1999;**72**:412–417.
- Nguyen Q, Sommer S, Greene B, Wrenzycki C, Wagner U, Ziller V. Effects of opening the incubator on morphokinetics in mouse embryos. *Eur J Obstet Gynecol Reprod Biol* 2018;**229**:64–69.
- Noli L, Dajani Y, Capalbo A, Bvumbe J, Rienzi L, Ubaldi FM, Ogilvie C, Khalaf Y, Ilic D. Developmental clock compromises human twin model created by embryo splitting. *Hum Reprod* 2015;**30**:2774–2784.
- Orevich LS, Watson K, Ong K, Korman I, Turner R, Shaker D, Liu Y. Morphometric and morphokinetic differences in the sperm- and oocyte-originated pronuclei of male and female human zygotes: a time-lapse study. *J Assist Reprod Genet* 2022;**39**:97–106.
- Otsuki J, Iwasaki T, Enatsu N, Katada Y, Furuhashi K, Shiotani M. Noninvasive embryo selection: kinetic analysis of female and male pronuclear development to predict embryo quality and potential to produce live birth. *Fertil Steril* 2019;**112**:874–881.
- Otsuki J, Iwasaki T, Katada Y, Tsutsumi Y, Tsuji Y, Furuhashi K, Kokeguchi S, Shiotani M. A higher incidence of cleavage failure in oocytes containing smooth endoplasmic reticulum clusters. *J Assist Reprod Genet* 2018;**35**:899–905.
- Otsuki J, Iwasaki T, Tsuji Y, Katada Y, Sato H, Tsutsumi Y, Hatano K, Furuhashi K, Matsumoto Y, Kokeguchi S et al Potential of zygotes to produce live births can be identified by the size of the male and female pronuclei just before their membranes break down. *Reprod Med Biol* 2017;**16**:200–205.
- Otsuki J, Nagai Y, Chiba K. Lipofuscin bodies in human oocytes as an indicator of oocyte quality. *J Assist Reprod Genet* 2007;**24**:263–270.
- Otsuki J, Okada A, Morimoto K, Nagai Y, Kubo H. The relationship between pregnancy outcome and smooth endoplasmic reticulum clusters in MII human oocytes. *Hum Reprod* 2004;**19**:1591–1597.
- Pan C, Zhang H. Embryological characteristics and clinical outcomes of oocytes with heterogeneous zona pellucida during assisted reproduction treatment: a retrospective study. *Med Sci Monit* 2020;**26**:e924316.
- Parriego M, Coll L, Carrasco B, Garcia S, Boada M, Polyzos NP, Vidal F, Veiga A. Blastocysts from partial compaction morulae are not defined by their early mistakes. *Reprod Biomed Online* 2024;**48**:103729.
- Payne D, Flaherty SP, Barry MF, Matthews CD. Preliminary observations on polar body extrusion and pronuclear formation in human oocytes using time-lapse video cinematography. *Hum Reprod* 1997;**12**:532–541.
- Payne D, Okuda A, Wakatsuki Y, Takeshita C, Iwata K, Shimura T, Yumoto K, Ueno Y, Flaherty S, Mio Y. Time-lapse recording identifies human blastocysts at risk of producing monozygotic twins. *Hum Reprod* 2007;**22**:I9–I10.
- Paz MV, Chiera M, Hovanyecz P, Cicaré J, Perfumo P, Domenech L, Ventura V. Blastocysts derived from OPN oocytes: genetic and clinical results. *JBRA Assist Reprod* 2020;**24**:143–146.
- Petersen BM, Boel M, Montag M, Gardner DK. Development of a generally applicable morphokinetic algorithm capable of predicting the implantation potential of embryos transferred on Day 3. *Hum Reprod* 2016;**31**:2231–2244.
- Petersen CG, Oliveira JB, Mauri AL, Massaro FC, Baruffi RL, Pontes A, Franco JG Jr. Relationship between visualization of meiotic spindle in human oocytes and ICSI outcomes: a meta-analysis. *Reprod Biomed Online* 2009;**18**:235–243.
- Pierson HE, Invik J, Meriano J, Pierson RA. A novel system for rapid conversion of Gardner embryo grades to linear scale numeric variables. *Reprod Biomed Online* 2023;**46**:808–818.
- Pons MC, Carrasco B, Parriego M, Boada M, González-Foruria I, Garcia S, Coroleu B, Barri PN, Veiga A. Deconstructing the myth of poor prognosis for fast-cleaving embryos on day 3. Is it time to change the consensus? *J Assist Reprod Genet* 2019;**36**:2299–2305.
- Pons MC, Carrasco B, Rives N, Delgado A, Martínez-Moro A, Martínez-Granados L, Rodríguez I, Cairó O, Cuevas-Saiz I; SIG Embryology of ASEBIR. Predicting the likelihood of live birth: an objective and user-friendly blastocyst grading system. *Reprod Biomed Online* 2023;**47**:103243.
- Practice Committee of the American Society for Reproductive Medicine. Blastocyst culture and transfer in clinically assisted reproduction: a committee opinion. *Fertility and Sterility* 2018;**110**:1246–1252.
- Practice Committee of Society for Assisted Reproductive Technology; Practice Committee of American Society for Reproductive Medicine. Elective single-embryo transfer. *Fertil Steril* 2012;**97**:835–842.
- Pribenszky C, Nilselid AM, Montag M. Time-lapse culture with morphokinetic embryo selection improves pregnancy and live birth chances and reduces early pregnancy loss: a meta-analysis. *Reprod Biomed Online* 2017;**35**:511–520. and *reproductive biology* 2018;**230**:96–102.
- Puissant F, Van Rysselberge M, Barlow P, Deweze J, Leroy F. Embryo scoring as a prognostic tool in IVF treatment. *Hum Reprod* 1987;**2**:705–708.

- Quinn P, Harlow GM. The effect of oxygen on the development of preimplantation mouse embryos in vitro. *J Exp Zool* 1978; **206**:73–80.
- Racowsky C, Stern JE, Gibbons WE, Behr B, Pomeroy KO, Biggers JD. National collection of embryo morphology data into Society for Assisted Reproductive Technology Clinic Outcomes Reporting System: associations among day 3 cell number, fragmentation and blastomere asymmetry, and live birth rate. *Fertil Steril* 2011; **95**:1985–1989.
- Raja EA, Bhattacharya S, Maheshwari A, McLernon DJ. A comparison of perinatal outcomes following fresh blastocyst or cleavage stage embryo transfer in singletons and twins and between singleton siblings. *Hum Reprod Open* 2023; **2023**:hoad003.
- Rama Raju GA, Prakash GJ, Krishna KM, Madan K. Meiotic spindle and zona pellucida characteristics as predictors of embryonic development: a preliminary study using PolScope imaging. *Reprod Biomed Online* 2007; **14**:166–174.
- Rattanachaiyanont M, Leader A, Léveillé MC. Lack of correlation between oocyte-corona-cumulus complex morphology and nuclear maturity of oocytes collected in stimulated cycles for intracytoplasmic sperm injection. *Fertil Steril* 1999; **71**:937–940.
- Regin M, Spits C, Sermon K. On the origins and fate of chromosomal abnormalities in human preimplantation embryos: an unsolved riddle. *Mol Hum Reprod* 2022; **28**:gaac011. <https://doi.org/10.1093/molehr/gaac011>
- Rhenman A, Berglund L, Brodin T, Olovsson M, Milton K, Hadziosmanovic N, Holte J. Which set of embryo variables is most predictive for live birth? A prospective study in 6252 single embryo transfers to construct an embryo score for the ranking and selection of embryos. *Hum Reprod* 2015; **30**:28–36.
- Rienzi L, Balaban B, Ebner T, Mandelbaum J. The oocyte. *Hum Reprod* 2012; **27** Suppl 1:i2–i21.
- Rienzi L, Capalbo A, Stoppa M, Romano S, Maggiulli R, Albricci L, Scarica C, Farcomeni A, Vajta G, Ubaldi FM. No evidence of association between blastocyst aneuploidy and morphokinetic assessment in a selected population of poor-prognosis patients: a longitudinal cohort study. *Reprod Biomed Online* 2015; **30**:57–66.
- Rienzi L, Cimadomo D, Delgado A, Minasi MG, Fabozzi G, Gallego RD, Stoppa M, Bellver J, Gianciani A, Esbert M et al Time of morulation and trophoctoderm quality are predictors of a live birth after euploid blastocyst transfer: a multicenter study. *Fertil Steril* 2019; **112**:1080–1093.e1081.
- Rienzi L, Ubaldi F, Minasi MG, Iacobelli M, Martinez F, Tesarik J, Greco E. Blastomere cytoplasmic granularity is unrelated to developmental potential of day 3 human embryos. *J Assist Reprod Genet* 2003; **20**:314–317.
- Rienzi L, Ubaldi FM, Iacobelli M, Minasi MG, Romano S, Ferrero S, Sapienza F, Baroni E, Litwicka K, Greco E. Significance of metaphase II human oocyte morphology on ICSI outcome. *Fertil Steril* 2008; **90**:1692–1700.
- Rienzi L, Vajta G, Ubaldi F. Predictive value of oocyte morphology in human IVF: a systematic review of the literature. *Hum Reprod Update* 2011; **17**:34–45.
- Rosenbusch B, Schneider M, Gläser B, Brucker C. Cytogenetic analysis of giant oocytes and zygotes to assess their relevance for the development of digynic triploidy. *Hum Reprod* 2002; **17**:2388–2393.
- Rubino P, Viganò P, Luddi A, Piomboni P. The ICSI procedure from past to future: a systematic review of the more controversial aspects. *Hum Reprod Update* 2016; **22**:194–227.
- Sá R, Cunha M, Silva J, Luís A, Oliveira C, Teixeira da Silva J, Barros A, Sousa M. Ultrastructure of tubular smooth endoplasmic reticulum aggregates in human metaphase II oocytes and clinical implications. *Fertil Steril* 2011; **96**:143–149.e147.
- Sacha CR, Kaser DJ, Farland LV, Srouji S, Missmer SA, Racowsky C. The effect of short-term exposure of cumulus-oocyte complexes to in vitro maturation medium on yield of mature oocytes and usable embryos in stimulated cycles. *J Assist Reprod Genet* 2018; **35**:841–849.
- Salas-Vidal E, Lomelí H. Imaging filopodia dynamics in the mouse blastocyst. *Dev Biol* 2004; **265**:75–89.
- Salumets A, Hydén-Granskog C, Mäkinen S, Suikkari AM, Tiitinen A, Tuuri T. Early cleavage predicts the viability of human embryos in elective single embryo transfer procedures. *Hum Reprod* 2003; **18**:821–825.
- Sathananthan AH. Ultrastructural changes during meiotic maturation in mammalian oocytes: unique aspects of the human oocyte. *Microsc Res Tech* 1994; **27**:145–164.
- Sauerbrun-Cutler MT, Vega M, Breborowicz A, Gonzales E, Stein D, Lederman M, Keltz M. Oocyte zona pellucida dysmorphology is associated with diminished in-vitro fertilization success. *J Ovarian Res* 2015; **8**:5.
- Savio Figueira Rde C, Setti AS, Braga DP, Iaconelli A Jr, Borges E Jr. Blastocyst morphology holds clues concerning the chromosomal status of the embryo. *Int J Fertil Steril* 2015; **9**:215–220.
- Sawada Y, Sato T, Nagaya M, Saito C, Yoshihara H, Banno C, Matsumoto Y, Matsuda Y, Yoshikai K, Sawada T et al Evaluation of artificial intelligence using time-lapse images of IVF embryos to predict live birth. *Reprod Biomed Online* 2021; **43**:843–852.
- Sayed S, Reigstad MM, Petersen BM, Schwennicke A, Hausken JW, Storeng R. Nucleation status of Day 2 pre-implantation embryos, acquired by time-lapse imaging during IVF, is associated with live birth. *PLoS One* 2022; **17**:e0274502.
- Sayed S, Reigstad MM, Petersen BM, Schwennicke A, Wegner Hausken J, Storeng R. Time-lapse imaging derived morphokinetic variables reveal association with implantation and live birth following in vitro fertilization: a retrospective study using data from transferred human embryos. *PLoS One* 2020; **15**:e0242377.
- Sciorio R, Herrero Saura R, Thong KJ, Esbert Algam M, Pickering SJ, Meseguer M. Blastocyst collapse as an embryo marker of low implantation potential: a time-lapse multicentre study. *Zygote* 2020; **28**:1–9.
- Scott KL, Hong KH, Scott RT Jr. Selecting the optimal time to perform biopsy for preimplantation genetic testing. *Fertil Steril* 2013; **100**:608–614.
- Scott L, Finn A, O'Leary T, McLellan S, Hill J. Morphologic parameters of early cleavage-stage embryos that correlate with fetal development and delivery: prospective and applied data for increased pregnancy rates. *Hum Reprod* 2007; **22**:230–240.
- Serdarogullari M, Findikli N, Goktas C, Sahin O, Ulug U, Yagmur E, Bahceci M. Comparison of gender-specific human embryo development characteristics by time-lapse technology. *Reprod Biomed Online* 2014; **29**:193–199.
- Serhal PF, Ranieri DM, Kinis A, Marchant S, Davies M, Khadum IM. Oocyte morphology predicts outcome of intracytoplasmic sperm injection. *Hum Reprod* 1997; **12**:1267–1270.
- Setti AS, Braga D, Vingris L, Iaconelli A Jr, Borges E Jr. Early and late paternal contribution to cell division of embryos in a time-lapse imaging incubation system. *Andrologia* 2021; **53**:e14211.
- Setti AS, Figueira RC, Braga DP, Colturato SS, Iaconelli A Jr, Borges E Jr. Relationship between oocyte abnormal morphology and intracytoplasmic sperm injection outcomes: a meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2011; **159**:364–370.
- Setti AS, Figueira RC, de Almeida Ferreira Braga DP, Azevedo MC, Iaconelli A Jr, Borges E Jr. Oocytes with smooth endoplasmic reticulum clusters originate blastocysts with impaired implantation potential. *Fertil Steril* 2016; **106**:1718–1724.

- Sfontouris IA, Lainas GT, Lainas TG, Faros E, Banti M, Kardara K, Anagnostopoulou K, Kontos H, Petsas GK, Kolibianakis EM. Complex chromosomal aberrations in a fetus originating from oocytes with smooth endoplasmic reticulum (SER) aggregates. *Syst Biol Reprod Med* 2018;**64**:283–290.
- Sha QQ, Zheng W, Wu YW, Li S, Guo L, Zhang S, Lin G, Ou XH, Fan HY. Dynamics and clinical relevance of maternal mRNA clearance during the oocyte-to-embryo transition in humans. *Nat Commun* 2020;**11**:4917.
- Shani AK, Haham LM, Balakier H, Kuznyetsova I, Bashar S, Day EN, Librach CL. The developmental potential of mature oocytes derived from rescue in vitro maturation. *Fertil Steril* 2023;**120**:860–869.
- Shaw-Jackson C, Thomas AL, Van Beirs N, Ameye L, Colin J, Bertrand E, Becker B, Rozenberg S, Autin C. Oocytes affected by smooth endoplasmic reticulum aggregates: to discard or not to discard? *Arch Gynecol Obstet* 2016;**294**:175–184.
- Shebl O, Haslinger C, Kresic S, Enengl S, Reiter E, Oppelt P, Ebner T. The hare and the tortoise: extreme mitotic rates and how these affect live birth. *Reprod Biomed Online* 2021;**42**:332–339.
- Shenoy CC, Khan Z, Coddington CC, Stewart EA, Morbeck DE. Symmetry at the 4-cell stage is associated with embryo aneuploidy. *Reprod Sci* 2021;**28**:3473–3479.
- Shi W, Xu B, Wu LM, Jin RT, Luan HB, Luo LH, Zhu Q, Johansson L, Liu YS, Tong XH. Oocytes with a dark zona pellucida demonstrate lower fertilization, implantation and clinical pregnancy rates in IVF/ICSI cycles. *PLoS One* 2014;**9**:e89409.
- Shoukir Y, Campana A, Farley T, Sakkas D. Early cleavage of in-vitro fertilized human embryos to the 2-cell stage: a novel indicator of embryo quality and viability. *Hum Reprod* 1997;**12**:1531–1536.
- Shu Y, Gebhardt J, Watt J, Lyon J, Dasig D, Behr B. Fertilization, embryo development, and clinical outcome of immature oocytes from stimulated intracytoplasmic sperm injection cycles. *Fertil Steril* 2007;**87**:1022–1027.
- Si J, Zhu X, Lyu Q, Kuang Y. Obstetrical and neonatal outcomes after transfer of cleavage-stage and blastocyst-stage embryos derived from monopronuclear zygotes: a retrospective cohort study. *Fertil Steril* 2019;**112**:527–533.
- Siristatidis C, Papapanou M, Karageorgiou V, Martins WP, Bellos I, Teixeira DM, Vlahos N. Congenital anomaly and perinatal outcome following blastocyst- vs cleavage-stage embryo transfer: systematic review and network meta-analysis. *Ultrasound Obstet Gynecol* 2023;**61**:12–25.
- Skiadas CC, Jackson KV, Racowsky C. Early compaction on day 3 may be associated with increased implantation potential. *Fertil Steril* 2006;**86**:1386–1391.
- Smith R, Barrie A, Berrisford K, Corcoran S, Davis N, Montgomery S, Oakley R, Owen S, Campbell A. Does the use of an in-house derived statistical morphokinetic algorithm, predictive of live birth, introduce any sex-ratio bias? A review of 3009 live birth outcomes. *Reproductive Biomedicine Online* 2024;**48**:104027.
- Sousa M, Cunha M, Silva J, Oliveira E, Pinho MJ, Almeida C, Sá R, da Silva JT, Oliveira C, Barros A. Ultrastructural and cytogenetic analyses of mature human oocyte dysmorphisms with respect to clinical outcomes. *J Assist Reprod Genet* 2016;**33**:1041–1057.
- Sousa M, Teixeira da Silva J, Silva J, Cunha M, Viana P, Oliveira E, Sá R, Soares C, Oliveira C, Barros A. Embryological, clinical and ultrastructural study of human oocytes presenting indented zona pellucida. *Zygote* 2015;**23**:145–157.
- Squirrel JM, Schramm RD, Paprocki AM, Wokosin DL, Bavister BD. Imaging mitochondrial organization in living primate oocytes and embryos using multiphoton microscopy. *Microsc Microanal* 2003;**9**:190–201.
- Storr A, Bilir E, Cooke S, Garrett D, Venetis CA. Fine-tuning blastocyst selection based on morphology: a multicentre analysis of 2461 single blastocyst transfers. *Reprod Biomed Online* 2019;**39**:588–598.
- Storr A, Venetis CA, Cooke S, Kilani S, Ledger W. Inter-observer and intra-observer agreement between embryologists during selection of a single Day 5 embryo for transfer: a multicenter study. *Hum Reprod* 2017;**32**:307–314.
- Strassburger D, Goldstein A, Friedler S, Raziel A, Kasterstein E, Mashevich M, Schachter M, Ron-El R, Reish O. The cytogenetic constitution of embryos derived from immature (metaphase I) oocytes obtained after ovarian hyperstimulation. *Fertil Steril* 2010;**94**:971–978.
- Subira J, Craig J, Turner K, Bevan A, Ohuma E, McVeigh E, Child T, Fatum M. Grade of the inner cell mass, but not trophectoderm, predicts live birth in fresh blastocyst single transfers. *Hum Fertil (Camb)* 2016;**19**:254–261.
- Sundström P, Saldeen P. Early embryo cleavage and day 2 mononucleation after intracytoplasmic sperm injection for predicting embryo implantation potential in single embryo transfer cycles. *Fertil Steril* 2008;**89**:475–477.
- Swain JE. Decisions for the IVF laboratory: comparative analysis of embryo culture incubators. *Reprod Biomed Online* 2014;**28**:535–547.
- Swain JE. Optimal human embryo culture. *Semin Reprod Med* 2015;**33**:103–117.
- Swain JE. Controversies in ART: considerations and risks for uninterrupted embryo culture. *Reprod Biomed Online* 2019;**39**:19–26.
- Tabibnejad N, Soleimani M, Aflatoonian A. Zona pellucida birefringence and meiotic spindle visualization are not related to the time-lapse detected embryo morphokinetics in women with polycystic ovarian syndrome. *Eur J Obstet Gynecol Reprod Biol* 2018;**230**:96–102.
- Takahashi H, Otsuki J, Yamamoto M, Saito H, Hirata REI, Habara T, Hayashi N. Clinical outcomes of MII oocytes with refractile bodies in patients undergoing ICSI and single frozen embryo transfer. *Reprod Med Biol* 2020;**19**:75–81.
- Talbot AL, Alexopoulou E, Kallemosse T, Freiesleben NC, Nielsen HS, Zedeler A. Binucleated embryos at the two-cell stage show higher blastocyst formation rates and higher pregnancy and live birth rates compared to non-multinucleated embryos. *Hum Reprod Open* 2022;**2022**:hoac049.
- Tannus S, Cohen Y, Henderson S, Al Ma'mari N, Shavit T, Son WY, Dahan MH. Fresh transfer of Day 5 slow-growing embryos versus deferred transfer of vitrified, fully expanded Day 6 blastocysts: which is the optimal approach? *Hum Reprod* 2019;**34**:44–51.
- Tao J, Tamis R, Fink K, Williams B, Nelson-White T, Craig R. The neglected morula/compact stage embryo transfer. *Hum Reprod* 2002;**17**:1513–1518.
- Ten J, Mendiola J, Vioque J, de Juan J, Bernabeu R. Donor oocyte dysmorphisms and their influence on fertilization and embryo quality. *Reprod Biomed Online* 2007;**14**:40–48.
- Tesarik J, Greco E. The probability of abnormal preimplantation development can be predicted by a single static observation on pronuclear stage morphology. *Hum Reprod* 1999;**14**:1318–1323.
- The Cairo Consensus Group. 'There is only one thing that is truly important in an IVF laboratory: everything' Cairo Consensus Guidelines on IVF Culture Conditions. *Reprod Biomed Online* 2020;**40**:33–60.
- Thompson SM, Onwubalili N, Brown K, Jindal SK, McGovern PG. Blastocyst expansion score and trophectoderm morphology strongly predict successful clinical pregnancy and live birth following elective single embryo blastocyst transfer (eSET): a national study. *J Assist Reprod Genet* 2013;**30**:1577–1581.

- Thurin A, Hardarson T, Hausken J, Jablonowska B, Lundin K, Pinborg A, Bergh C. Predictors of ongoing implantation in IVF in a good prognosis group of patients. *Hum Reprod* 2005;**20**:1876–1880.
- Tian L, Xia L, Liu H, Kou YAN, Huang Z, Wu X, Fan LU, Huang J, Wu Q. Increased blastomere number is associated with higher live birth rate in day 3 embryo transfer. *BMC Pregnancy Childbirth* 2022;**22**:198.
- Tiegs AW, Sun L, Patounakis G, Scott RT. Worth the wait? Day 7 blastocysts have lower euploidy rates but similar sustained implantation rates as Day 5 and Day 6 blastocysts. *Hum Reprod* 2019;**34**:1632–1639.
- Tiegs AW, Tao X, Zhan Y, Whitehead C, Kim J, Hanson B, Osman E, Kim TJ, Patounakis G, Gutmann J et al A multicenter, prospective, blinded, nonselection study evaluating the predictive value of an aneuploid diagnosis using a targeted next-generation sequencing-based preimplantation genetic testing for aneuploidy assay and impact of biopsy. *Fertil Steril* 2021;**115**:627–637.
- Tran D, Cooke S, Illingworth PJ, Gardner DK. Deep learning as a predictive tool for fetal heart pregnancy following time-lapse incubation and blastocyst transfer. *Hum Reprod* 2019;**34**:1011–1018.
- Ubbaldi FM, Capalbo A, Vaiarelli A, Cimadomo D, Colamaria S, Alviggi C, Trabucco E, Venturella R, Vajta G, Rienzi L. Follicular versus luteal phase ovarian stimulation during the same menstrual cycle (DuoStim) in a reduced ovarian reserve population results in a similar euploid blastocyst formation rate: new insight in ovarian reserve exploitation. *Fertil Steril* 2016;**105**:1488–1495.e1481.
- Ueno S, Bodri D, Uchiyama K, Okimura T, Okuno T, Kobayashi T, Kato K. Developmental potential of zona pellucida-free oocytes obtained following mild in vitro fertilization. *Fertil Steril* 2014;**102**:1602–1607.
- Van de Velde H, De Vos A, Sermon K, Staessen C, De Rycke M, Van Assche E, Lissens W, Vandervorst M, Van Ranst H, Liebaers I et al Embryo implantation after biopsy of one or two cells from cleavage-stage embryos with a view to preimplantation genetic diagnosis. *Prenat Diagn* 2000;**20**:1030–1037.
- Van den Abbeel E, Balaban B, Ziebe S, Lundin K, Cuesta MJ, Klein BM, Helmggaard L, Arce JC. Association between blastocyst morphology and outcome of single-blastocyst transfer. *Reprod Biomed Online* 2013;**27**:353–361.
- Van Montfoort AP, Dumoulin JC, Kester AD, Evers JL. Early cleavage is a valuable addition to existing embryo selection parameters: a study using single embryo transfers. *Hum Reprod* 2004;**19**:2103–2108.
- Van Royen E, Mangelschots K, De Neubourg D, Laureys I, Ryckaert G, Gerris J. Calculating the implantation potential of day 3 embryos in women younger than 38 years of age: a new model. *Hum Reprod* 2001;**16**:326–332.
- Van Royen E, Mangelschots K, Vercruyssen M, De Neubourg D, Valkenburg M, Ryckaert G, Gerris J. Multinucleation in cleavage stage embryos. *Hum Reprod* 2003;**18**:1062–1069.
- Vassena R, Boué S, González-Roca E, Aran B, Auer H, Veiga A, Izpisua Belmonte JC. Waves of early transcriptional activation and pluripotency program initiation during human preimplantation development. *Development* 2011;**138**:3699–3709.
- Veeck LL, Zaninovic N. *An Atlas of Human Blastocysts*. Boca Raton, USA: CRC Press, 2003.
- Vera-Rodriguez M, Chavez SL, Rubio C, Reijo Pera RA, Simon C. Prediction model for aneuploidy in early human embryo development revealed by single-cell analysis. *Nat Commun* 2015;**6**:7601.
- Verlinsky Y, Lerner S, Illkevitch N, Kuznetsov V, Kuznetsov I, Cieslak J, Kuliev A. Is there any predictive value of first polar body morphology for embryo genotype or developmental potential? *Reprod Biomed Online* 2003;**7**:336–341.
- Vitthala S, Gelbaya TA, Brison DR, Fitzgerald CT, Nardo LG. The risk of monozygotic twins after assisted reproductive technology: a systematic review and meta-analysis. *Hum Reprod Update* 2009;**15**:45–55.
- Wale PL, Gardner DK. Time-lapse analysis of mouse embryo development in oxygen gradients. *Reprod Biomed Online* 2010;**21**:402–410.
- Wale PL, Gardner DK. The effects of chemical and physical factors on mammalian embryo culture and their importance for the practice of assisted human reproduction. *Hum Reprod Update* 2016;**22**:2–22.
- Wallbuton S, Kasraie J. Vacuolated oocytes: fertilization and embryonic arrest following intra-cytoplasmic sperm injection in a patient exhibiting persistent oocyte macro vacuolization—case report. *J Assist Reprod Genet* 2010;**27**:183–188.
- Wang M, Gao L, Yang Q, Long R, Zhang Y, Jin L, Zhu L. Does smooth endoplasmic reticulum aggregation in oocytes impact the chromosome aneuploidy of the subsequent embryos? A propensity score matching study. *J Ovarian Res* 2023;**16**:59.
- Wang Q, Ulker A, Wang H, Wu B, Yang A, Attia GR. Single live birth derived from conjoined oocytes using laser-cutting technique: a case report. *Zygote* 2022;**30**:217–220.
- Wang X, Du M, Guan Y, Wang B, Zhang J, Liu Z. Comparative neonatal outcomes in singleton births from blastocyst transfers or cleavage-stage embryo transfers: a systematic review and meta-analysis. *Reprod Biol Endocrinol* 2017;**15**:36.
- Wang X, Xiao Y, Sun Z, Zhen J, Yu Q. Smooth endoplasmic reticulum clusters in oocytes from patients who received intracytoplasmic sperm injections negatively affect blastocyst quality and speed of blastocyst development. *Front Physiol* 2021;**12**:732547.
- Weghofer A, Kushnir VA, Darmon SK, Jafri H, Lazzaroni-Tealdi E, Zhang L, Albertini DF, Barad DH, Gleicher N. Age, body weight and ovarian function affect oocyte size and morphology in non-PCOS patients undergoing intracytoplasmic sperm injection (ICSI). *PLoS One* 2019;**14**:e0222390.
- Wei X, Enatsu N, Furuhashi K, Iwasaki T, Kokeguchi S, Shiotani M, Otsuki J. Developmental trajectory of mononucleated zygotes after in vitro fertilization when they include both male and female genomes. *Fertil Steril* 2022;**117**:213–220.
- White CR, Denomme MM, Tekpetey FR, Feyles V, Power SG, Mann MR. High frequency of imprinted methylation errors in human preimplantation embryos. *Sci Rep* 2015;**5**:17311.
- Wilding M, Di Matteo L, D'Andretti S, Montanaro N, Capobianco C, Dale B. An oocyte score for use in assisted reproduction. *J Assist Reprod Genet* 2007;**24**:350–358.
- Wu J, Zhang J, Kuang Y, Chen Q, Wang Y. The effect of Day 3 cell number on pregnancy outcomes in vitrified-thawed single blastocyst transfer cycles. *Hum Reprod* 2020;**35**:2478–2487.
- Wyns C, De Geyter C, Calhaz-Jorge C, Kupka MS, Motrenko T, Smeenk J, Bergh C, Tandler-Schneider A, Rugesu IA, Goossens V; European IVF Monitoring Consortium (EIM), for the European Society of Human Reproduction and Embryology (ESHRE). ART in Europe, 2018: results generated from European registries by ESHRE. *Hum Reprod Open* 2022;**2022**:hoac022.
- Xia P. Intracytoplasmic sperm injection: correlation of oocyte grade based on polar body, perivitelline space and cytoplasmic inclusions with fertilization rate and embryo quality. *Hum Reprod* 1997;**12**:1750–1755.
- Xie PY, Tang Y, Hu L, Ouyang Q, Gu YF, Gong F, Leng LZ, Zhang SP, Xiong B, Lu GX et al Identification of biparental and diploid blastocysts from monopronuclear zygotes with the use of a single-nucleotide polymorphism array. *Fertil Steril* 2018;**110**:545–554.e5.

- Xu J, Yang L, Chen ZH, Yin MN, Chen J, Sun L. Oocytes with smooth endoplasmic reticulum aggregates may not impact blastocyst euploidy rate. *Front Endocrinol (Lausanne)* 2022;**13**:851370.
- Yakin K, Balaban B, Isiklar A, Urman B. Oocyte dysmorphism is not associated with aneuploidy in the developing embryo. *Fertil Steril* 2007;**88**:811–816.
- Yalçınkaya E, Özay A, Ergin EG, Öztel Z, Özörnek H. Live birth after transfer of a tripronuclear embryo: An intracytoplasmic sperm injection as a combination of microarray and time-lapse technology. *Turk J Obstet Gynecol* 2016;**13**:95–98.
- Yan L, Huang L, Xu L, Huang J, Ma F, Zhu X, Tang Y, Liu M, Lian Y, Liu P et al Live births after simultaneous avoidance of monogenic diseases and chromosome abnormality by next-generation sequencing with linkage analyses. *Proc Natl Acad Sci USA* 2015;**112**:15964–15969.
- Yang D, Yang H, Yang B, Wang K, Zhu Q, Wang J, Ding F, Rao B, Xue R, Peng J et al Embryological characteristics of human oocytes with agar-like zona pellucida and its clinical treatment strategy. *Front Endocrinol (Lausanne)* 2022;**13**:859361.
- Yang Q, Zhu L, Wang M, Huang B, Li Z, Hu J, Xi Q, Liu J, Jin L. Analysis of maturation dynamics and developmental competence of in vitro matured oocytes under time-lapse monitoring. *Reprod Biol Endocrinol* 2021;**19**:183.
- Yerushalmi GM, Shavit T, Avraham S, Youngster M, Kedem A, Gat I, Dorofeyeva US, Mashiach S, Schiff E, Shulman A et al. Day 5 vitrified blastocyst transfer versus day 6 vitrified blastocyst transfer in oocyte donation program. *Sci Rep* 2021;**11**:10715.
- Yang ST, Shi JX, Gong F, Zhang SP, Lu CF, Tan K, Leng LZ, Hao M, He H, Gu YF et al Cleavage pattern predicts developmental potential of day 3 human embryos produced by IVF. *Reprod Biomed Online* 2015;**30**:625–634.
- Yang Z, Zhang J, Salem SA, Liu X, Kuang Y, Salem RD, Liu J. Selection of competent blastocysts for transfer by combining time-lapse monitoring and array CGH testing for patients undergoing preimplantation genetic screening: a prospective study with sibling oocytes. *BMC Med Genomics* 2014;**7**:38.
- Yi XF, Xi HL, Zhang SL, Yang J. Relationship between the positions of cytoplasmic granulation and the oocytes developmental potential in human. *Sci Rep* 2019;**9**:7215.
- Yu CH, Zhang RP, Li J, A ZC. A predictive model for high-quality blastocyst based on blastomere number, fragmentation, and symmetry. *J Assist Reprod Genet* 2018;**35**:809–816.
- Yuan S, Zhan J, Zhang J, Liu Z, Hou Z, Zhang C, Yi L, Gao L, Zhao H, Chen ZJ et al Human zygotic genome activation is initiated from paternal genome. *Cell Discov* 2023;**9**:13.
- Zhan Q, Sierra ET, Malmsten J, Ye Z, Rosenwaks Z, Zaninovic N. Blastocyst score, a blastocyst quality ranking tool, is a predictor of blastocyst ploidy and implantation potential. *F S Rep* 2020;**1**:133–141.
- Zhan Q, Ye Z, Clarke R, Rosenwaks Z, Zaninovic N. Direct unequal cleavages: embryo developmental competence, genetic constitution and clinical outcome. *PLoS One* 2016;**11**:e0166398.
- Zhang H, Hu W, Zhong Y, Guo Z. Meta-analysis of the effects of smooth endoplasmic reticulum aggregation on birth outcome. *BMC Pregnancy Childbirth* 2021;**21**:374.
- Zhang JQ, Li XL, Peng Y, Guo X, Heng BC, Tong GQ. Reduction in exposure of human embryos outside the incubator enhances embryo quality and blastulation rate. *Reprod Biomed Online* 2010;**20**:510–515.
- Zhang WY, Johal JK, Gardner RM, Bavan B, Milki AA. The impact of euploid blastocyst morphology and maternal age on pregnancy and neonatal outcomes in natural cycle frozen embryo transfers. *J Assist Reprod Genet* 2022;**39**:647–654.
- Zhao H, Liu H, Li M, Wu K. Clinical outcomes following frozen-thawed blastocyst transfers with blastocysts derived from different cell numbers on day 3: a retrospective cohort study. *J Assist Reprod Genet* 2020;**37**:641–648.
- Zhao H, Yuan P, Chen X, Lin H, Zhao J, Huang J, Qiu Q, Ji X, Zhang Q, Wang W. The aneuploidy testing of blastocysts developing from OPN and 1PN zygotes in conventional IVF through TE-biopsy PGT-A and minimally invasive PGT-A. *Front Reprod Health* 2022;**4**:966909.
- Zhao YY, Yu Y, Zhang XW. Overall blastocyst quality, trophoctoderm grade, and inner cell mass grade predict pregnancy outcome in euploid blastocyst transfer cycles. *Chin Med J (Engl)* 2018;**131**:1261–1267.
- Zhou W, Fu L, Sha W, Chu D, Li Y. Relationship of polar bodies morphology to embryo quality and pregnancy outcome. *Zygote* 2016;**24**:401–407.
- Zhu M, Shahbazi M, Martin A, Zhang C, Sozen B, Borsos M, Mandelbaum RS, Paulson RJ, Mole MA, Esbert M et al Human embryo polarization requires PLC signaling to mediate trophoctoderm specification. *Elife* 2021;**10**:e65068.
- Ziebe S. Morphometric analysis of human embryos to predict developmental competence. *Reprod Fertil Dev* 2013;**26**:55–64.
- Ziebe S, Lundin K, Loft A, Bergh C, Nyboe Andersen A, Selleskog U, Nielsen D, Grøndahl C, Kim H, Arce JC; CEMAS II and Study Group. FISH analysis for chromosomes 13, 16, 18, 21, 22, X and Y in all blastomeres of IVF pre-embryos from 144 randomly selected donated human oocytes and impact on pre-embryo morphology. *Hum Reprod* 2003;**18**:2575–2581.
- Ziebe S, Petersen K, Lindenberg S, Andersen AG, Gabrielsen A, Andersen AN. Embryo morphology or cleavage stage: how to select the best embryos for transfer after in-vitro fertilization. *Hum Reprod* 1997;**12**:1545–1549.
- Zou H, Kemper JM, Hammond ER, Xu F, Liu G, Xue L, Bai X, Liao H, Xue S, Zhao S et al Blastocyst quality and reproductive and perinatal outcomes: a multinational multicentre observational study. *Hum Reprod* 2023;**38**:2391–2399.
- Zou Y, Pan Y, Ge N, Xu Y, Gu R, Li Z, Fu J, Gao J, Sun X, Sun Y. Can the combination of time-lapse parameters and clinical features predict embryonic ploidy status or implantation? *Reprod Biomed Online* 2022;**45**:643–651.